

REMARKS

The Office Action of May 14, 2004 presents the Examination of claims 1-10, 16-23 and 28-30. These claims are not amended and remain pending.

Substitute Specification Provided

The Examiner objects to the specification as presenting an overly detailed Summary of the Invention portion that amounts to merely a restatement of the language of the claims. The Examiner's objection is addressed by provision of a substitute specification deleting that portion of the specification.

The Examiner also objects to informalities of a specification "filed April 14, 2000," related to incongruity between the description of sequences presented in the Sequence Listing and sequence listing identifier numbering in the instant specification. It appears that this objection arises due to confusion by the Examiner of the specification in the instant application and the specification of record in the co-pending application 08/992,914. Applicants note first that there was no specification filed April 14, 2000 in the present application. Applicants further note that the present specification does not include any Example 10 or 11 and furthermore that the Example 6 appears on page 45, not on page 37. Thus, it appears that the Examiner's reasons for objecting to the specification are not correct.

Applicants have incorporated the amendments of the Sequence Listing identifiers from the Preliminary Amendment of April 29, 1999 into the attached substitute specification.

No new matter is added by the substitute specification. A "track-changes" version of the substitute specification is attached hereto for the Examiner's review.

Rejection under 35 U.S.C. § 112, second paragraph

Claims 1-10, 16-23 and 28-30 are rejected under 35 U.S.C. § 112, second paragraph, for allegedly being indefinite in the recitation of incorrect Sequence Listing identifier numbers. Applicants believe that the Examiner bases this rejection upon reading of the wrong specification and therefore the rejection should be withdrawn.

Rejection under 35 U.S.C. § 101 and 35 U.S.C. § 112, first paragraph

Claims 1-10, 16-23 and 28-30 stand rejected under 35 U.S.C. § 101 and 35 U.S.C. § 112, first paragraph, for alleged lack of description of a utility or assertion of a known utility for the claimed invention. Claims 1-10, 16-23 and 28-30 are also rejected under 35 U.S.C. § 112, first paragraph for alleged lack of adequate written description of the invention. These rejections are respectfully traversed. Reconsideration and withdrawal thereof are requested.

The Examiner takes a position that there is no evidence of record that the claimed nucleic acids encode a protein having raffinose synthase activity. The Examiner asserts that mere homology to a known raffinose synthase gene is not sufficient, as there are other genes known that have higher degrees of homology to some raffinose synthase gene, yet encode proteins having other activities. In particular, the Examiner has compared the *S. affinis* stachyose synthase to the *S. cuminis* raffinose synthase and found a degree of homology of 50% and with the *P. sativum* raffinose synthase and found a degree of homology of 51%. Applicants note that in their prior response, a sequence comparison was provided as Table 2; these same comparisons are in Table 2 and a degree of homology of only 43% is found for both.

The discrepancy in the degree of "homology" in the data set provided by the Applicant and by the Examiner is due to differences in the computer program used to analyze the data. Table 2 of Applicants' previous response shows overall sequence homologies(%) between raffinose synthases (RFSs), imbibition protein (SIP) and stachyose synthases (STSs). Applicants' homology data were calculated using Global Alignment (the alignment of sequences over their entire length) produced by the CLUSTAL sequence analysis program. The CLUSTAL program uses the algorithm of Wilbur and Lipman (see the attached Exhibit 1, a description of sequence

analysis programs found at

http://www.rfcgr.mrc.ac.uk/embnet.news/vol2_1/align.html).

On the other hand, the Examiner's analysis was conducted using the BLAST program, which is a Basic Local Alignment Search Tool. The BLAST program uses the BLAST algorithm and makes local alignments (the alignment of some portion of two sequences) in order to search similarities of sequences. BLAST is a local alignment program, and does not make global alignments between sequences to calculate total percent homologies. (See, page 3, the 9th paragraph of attached Exhibit 2, http://www.ncbi.nlm.nih.gov/BLAST/blast_FAQs.shtml). Moreover, "identities" values output in a BLAST search report are different from "homology" values. That is, "homology" means "similarity attributed to descent from a common ancestor", while "identity" means "the extent to which two sequences are invariant" (See, pages 2-3 of attached Exhibit 3, <http://www.ncbi.nlm.nih.gov/Education/BLASTinfo/glossary2.html>). Thus, different values and scores calculated using different algorithms are based on different standards or criteria, and therefore it is not reasonable to discuss homology and similarity of sequences only by directly comparing such different values and scores.

The attached Table 3 shows the identities obtained using a similarity search using the BLAST program for the amino acid sequences of RFSSs, SIP and STSS shown in Table 1 (attached to

Applicants' previous response). Among Sc-02, Sc-03, Sc-04 and Sc-05, the identities were obtained by searching a patent database with default parameters, using the amino acid sequence of each protein as the "query", and using "Protein query vs. translated database (tblastn)" of the NCBI BLAST program. Also, other identities were obtained by searching the non-redundant database with default parameters, using the amino acid sequence of each protein as the "query", and using "Protein-protein BLAST (blastp)" of the NCBI BLAST program.

The identities between RFSSs and SIPs are about 40%. The identities between RFSSs and STSSs range from about 40% to about 50%. On the other hand, the identities among RFSSs are 60% or more. The identities among STSSs are also 60% or more. That is, the identities among RFSSs or the identities among STSSs are higher than the identities between RFSSs and SIPs or the identities between RFSSs and STSSs. Thus, based on the results of analyses by BLAST program, RFSSs, SIPs or STSSs can be distinguished. Applicants note that the conclusion reached from this analysis is consistent with the conclusion reached using the CLUSTAL analysis provided in their previous paper.

Applicants have also previously provided the Declaration of Dr. Watanabe to address the issues of adequacy of the written description and disclosure of utility. The Examiner has dismissed the Watanabe Declaration, stating that it is unclear what nucleic acid was used in the examples in the Declaration. The Watanabe

Declaration describes on its page 2, at the beginning of the item "Experiments":

The vector BjRS-Sac(+)-121 having the mustard raffinose synthase gene of the present invention in the expressible direction (i.e. sense direction) and the vector BjRS-Sac(-)-121 having the mustard raffinose synthase gene of the present invention in the reverse direction (i.e. antisense direction), which are the same as obtained in Example 8 of the present specification, were used for the transformation of tobacco (*Nicotiana tubacum*).

The Examiner should take note that Example 8 of the originally filed specification (page 47, lines 13-16) describes "the nucleotide sequence of SEQ ID NO:6 and SEQ ID NO:8 were determined from mustard (*Brassica Juncea*) and rapeseed Wester (*Brassica napus*), respectively". Therefore, the mustard raffinose synthase gene used in Watanabe Declaration is the gene from the mustard *Brassica Juncea*.

The Examiner also dismisses the Watanabe Declaration as presenting only a single example of a functional raffinose synthase gene. The Examiner's argument that Dr. Watanabe establishes functionality of only one gene is not persuasive. Dr. Watanabe's Declaration provides evidence of functionality in support of an assertion of utility that cannot be dismissed merely because of the number of examples it sets forth. Dr. Watanabe's Declaration provides evidence that the asserted utility for the invention is indeed correct; any "rebuttal" by the Examiner must be supported by

evidence, not a mere assertion that not enough examples have been provided.

The Examiner "rebutts" Applicants' arguments about the sufficiency of the disclosure in the specification by suggesting that the sequences mentioned in Applicants' arguments are not those set forth in the sequence listing. Applicants note the Examiner's confusion of the instant application with a co-pending application and urge that he revisit Applicants' previous argument.

The Examiner should note that it is Applicants' burden to establish adequacy of written description and utility only by the preponderance of the evidence. Applicants submit that the evidence of record firmly establishes 1) that raffinose synthase genes as a genus are more closely related to each other than to stachyose synthase genes or imbibition protein genes suggested by the Examiner as establishing lack of utility, and therefore the skilled artisan can distinguish a raffinose synthase gene from either of a stachyose synthase gene or an imbibition protein gene by proper homology analysis and 2) that a gene identified as a raffinose synthase gene by homology analysis is most likely to actually have the biochemical activity of a raffinose synthase gene. (This latter being established by the Watanabe Declaration at least.)

Accordingly, the utility of the instantly claimed invention is established and the rejection of claims 1-10, 16-23 and 28-30 stand rejected under 35 U.S.C. §§ 101 and 112, first paragraph,

for alleged lack of proof of an asserted utility should be withdrawn.

The adequacy of the written description of the invention is established by the comparison of the specification with the Guidelines for examination of written description in Applicants' previous response. The Examiner is requested to reconsider his position and arguments upon reading of Applicants' arguments together with the specification of the present application, not the specification of the co-pending application.

Obviousness-type double patenting

Claims 1-3, 16-23 and 28-30 are rejected under the judicially created doctrine of obviousness-type double patenting over claims of the co-pending application "08/992,766". First, the Examiner is requested to confirm that the co-pending application is in fact 08/992,914. Second, the Examiner is requested to hold this rejection in abeyance until either this application or the '914 application is allowed, at which time an appropriate response in the form of either arguments distinguishing the invention or a terminal disclaimer will be filed in the application that remains under examination.

Conclusion

Applicants respectfully submit that the above remarks and/or amendments fully address and overcome the rejections of record.

The present application is in condition for allowance. The Examiner is respectfully requested to issue a Notice of Allowance indicating that claims 1-10, 16-23, and 28-30 are allowed.


Should there be any outstanding matters that need to be resolved in the present application, the Examiner is respectfully requested to contact Mark J. Nuell, Ph.D. (Reg. No. 36,623) at the below-listed number.

Pursuant to the provisions of 37 C.F.R. §§ 1.17 and 1.136(a), the Applicants hereby petition for an extension of three (3) months to November 14, 2004, in which to file a reply to the Office Action. The required fee of \$980.00 is enclosed herewith.

If necessary, the Commissioner is hereby authorized in this, concurrent, and future replies, to charge payment or credit any overpayment to Deposit Account No. 02-2448 for any additional fees are required under 37 C.F.R. § 1.16 or under 37 C.F.R. § 1.17; particularly, extension of time fees.

Respectfully submitted,

BIRCH, STEWART, KOLASCH & BIRCH, LLP

By 
Mark J. Nuell, Reg. No. 36,623

GMM/DRN: kdm
0020-4559P

P.O. Box 747
Falls Church, VA 22040-0747
(703) 205-8000

Attachments: 1. Table 3: Homologies (%) among RFS, SIP, and STS enzymes;
2. Exhibits 1-3;
3. Substitute specification (track changes copy) and
4. Substitute specification (clean copy)

Table 3

[illegible]



ALGORITHMS FOR MULTIPLE SEQUENCE ALIGNMENTS

Guy Bottu,

BEN, The Belgian EMBnet node.

Introduction.

In a previous issue of embnetnews, we considered the alignment of pairs of sequences and the search for similar sequences in databanks. We now turn our attention to multiple sequence alignments.

If you have several similar nucleic acid or protein sequences it is often useful to align corresponding bases or amino acids in columns. For instance, you might wish to group bases or amino acids that occupy similar positions in the three-dimensional structure which exercise similar functions or that have evolved by substitution from the same base or amino acid in an ancestral sequence. In the latter case you might also like to construct a phylogenetic tree.

1. Global alignments.

The Needleman and Wunsch algorithm for finding the best global alignment of two sequences can readily be extended to multiple sequences. The problem is that the time the computer needs for such a job is roughly proportional to the product of the sequence lengths. So, if aligning two sequences of 300 positions takes 1 second, aligning 3 sequences takes 300 seconds and aligning 10 sequences would take 300^2 seconds, which is longer than the lifetime of the universe!

Since searching for a best global alignment using a rigorous algorithm is not realistic for more than three sequences, a number of strategies have been developed to carry out a multiple global alignment in a reasonable amount of time with a reasonable chance of finding the best alignment. The GCG program pileup first aligns all possible pairs of sequences according to Needleman and Wunsch (for n sequences, this makes $n(n-1)/2$ alignments). Then it uses the pairwise similarity scores to construct a tree using the UPGMA method (see below). Finally, this tree serves as a guide for a progressive multiple alignment starting from the tips. Once two sequences have been aligned, their relative alignment is no longer changed. Clusters of previously aligned sequences are treated as a linearly weighted profile when they are subsequently aligned with another sequence or another cluster.

Other approaches include:

- The very popular CLUSTAL program differs only from pileup in that it performs the initial pairwise alignments using the fast algorithm of Wilbur and Lipman. CABIOS 8:189 (1992). References
you can obtain versions of CLUSTAL for UNIX and for VAX
- Starting with a search for words of n bases or amino acids that are common between the sequences. An example is Martin Vingron's program MALI. CABIOS 5:115 (1989). References.
MALI is not distributed freely but may be obtained from its author Martin Vingron (vingron@embl-heidelberg.de)
- PIMA uses pattern-matching, rather than profile matching, while making the progressive alignment. PNAS 87:118 (1990) References
PIMA can be obtained for UNIX and for VAX
- Building a phylogenetic tree, using a more elaborate algorithm, as the sequences are progressively aligned. An example is Jotun Hein's program TreeAlign. Meth.Enzymol. 18:626(1990)
TreeAlign can be obtained for UNIX and VMS from the same address as given for Clustal (see above)
- Making the best multiple alignment in a limited area of alignment space. This can only realistically be performed with eight to ten sequences.

2. Local alignments.

There are cases where sequences share a similar region but are otherwise completely different. Take, for example, the amino acids in the active site of an enzyme or transcription factor binding sites in a DNA sequence. To handle these cases local multiple alignment algorithms have been developed. Usually they only look for ungapped alignments thereby avoiding the problem of choosing the optimal gap penalty. Two such programs have been developed at the NCBI :
MACAW by Schuler, Altschul and Lipman first tries to find high scoring segment pairs (HSPs) for each possible pair of sequences using the BLAST algorithm (with the sensitivity set high). It then assembles overlapping HSPs into blocks. An interesting feature of MACAW is that it does not try to align all sequences, but can pick out only those that share similar regions. Proteins 9:180 (1991). References

There are versions of MACAW obtainable for the PC under Windows and for the Mac.
The MACAW distribution also contains Gibbs (see below) and a pattern searcher.

The Gibbs sampler algorithm involves iteratively making a profile with stretches of n bases or amino acids, selected from the sequences, and then searches this profile against one of the sequences. The result of the search is used to weight the selection of the stretches at the next run. A drawback is that the user must choose the width n and the number of elements in each sequence and thus must have a certain idea of the outcome, or run the program several times. An interesting feature is that the Gibbs sampler algorithm avoids the choice of an externally added scoring scheme since it derives the highest scoring profile, in a

self-consistent manner, from the data. Science 262:208 (1992). References.
Gibbs for is available for UNIX.

3. blast3.

It is also worth mentioning the program blast3. This searches a protein against a protein databank using the BLAST algorithm (with the sensitivity set high) and then makes threefold alignments between the query sequence and each possible pair of databank sequences that have been found. Only the statistically significant threefold alignments which are made from three nonsignificant pairwise alignments are retained. blast3 is useful in finding proteins that share a region of only weak similarity. Occasionally it can show that a query sequence makes the bridge between two databank sequences whose relationship had not yet been suspected.

You can look at the Manual.

It is possible to access a BLAST (including blast3) server at the NCBI, either through WWW or with a specific blast Internet client that you can install on your computer. More INFO is available.

4. phylogenetic trees.

Ideally a researcher would like to have a black box in which to throw sequences and get out a fully annotated phylogenetic tree. This is, however, not possible for two reasons. First, an algorithm that considers all possible multiple sequence alignments and then, for each alignment, all possible phylogenetic trees and picks out the best one, would take too much time. That is why most phylogenetic programs work on previously aligned sequences. Second, the result is always strongly influenced by the criteria that are used to define the best tree. Phylogenetic analysis will be the subject of a separate column in a later issue of embnet.news. However, a few remarks seem appropriate here. There are three main kinds of tree building methods: distance matrix, maximum likelihood and parsimony.

Distance matrix methods first estimate the pairwise distances between the sequences (which means that the information in the alignment of two sequences is reduced to one number) while the other methods construct many trees from all the information in the multiple alignment and decide which is best.

The simplest distance based method is UPGMA (unweighted pair-group method using arithmetic averages) which involves iteratively taking together the two sequences that have the shortest distance from each other, placing them at the end of branches on a node of the tree, and replacing their distances from the other sequences by an average value.

The guide tree used by pileup and CLUSTAL should never be used to infer phylogeny! It has been derived from the distances between pairwise aligned sequences and these distances are not necessarily the same as the distances between sequence pairs taken from the multiple sequence alignment.

Go to: [previous article](#) - [next article](#) - [Table to contents](#)

**BLAST**PubMed
Info

Entrez

BLAST

OMIM

Taxonomy

Structure

BLAST Frequently Asked Questions (FAQ)Tips and Hints:

- FAQs
- News
- References
- NCBI Contributors

Education

- Program selection guide
- Tutorial
- URL API guide

Download

- Databases
- Documentation
- Executables
- Source code

Support

- Helpdesk
- Mailing list

- Which BLAST program should I use?
- How can I search a batch of sequences with BLAST?
- How can I write a program to submit jobs to NCBI's BLAST servers?
- How can I limit my BLAST search based on Organism?
- How can I limit my search to a subset of database sequences?
- Is it possible to search for a motif or pattern with BLAST?
- How do I perform a similarity search with a short peptide/nucleotide sequence?
- Can I use BLAST to compare two or more sequences in a multiple sequence alignment?
- What is the Expect (E) value?
- What is low-complexity sequence?
- Other Molecular Biology Resources

Troubleshooting:

- Why do I get the "No Significant similarity found" error message?
- Why does my search timeout on the BLAST servers?
- Why do I get the error message "ERROR: BLASTSetUpSearch: Unable to calculate Karlin-Altschul params, check query sequence"?
- Why do I get the error message "ERROR: Blast: No Valid Letters to be indexed?"
- Why do I see a string of "X"s (or "N") in my query sequence that I did not put there?
- I have heard that I will be penalized if I send a large number of sequences to the servers?

Tips and hints

Q: Which BLAST program should I use?

You have many choices to make between different BLAST programs and databases. Some of these choices are better for answering some questions than others. We have created a selection chart to help you make the decision of BLAST program for the question you are asking. This is the "BLAST Program Selection Guide".

Q: How can I search a batch of sequences with BLAST?

There are three options for "Batch" BLAST searches:

1) Web MegaBLAST EST analysis tool: This program is optimized for aligning nucleotide sequences that differ slightly as a result of sequencing or other similar "errors". MegaBLAST is good for scanning a large number of EST type sequences (about 500 kb in length) against large database in search of the closest matches. You can import a file EST sequences in FASTA format or as a list of GenBank accessions or/GIs and have them compared to the BLAST databases. The default is an easily reviewable Hit Table format, although you can download and save the results in Standard pairwise HTML or any of the other result output options. MegaBLAST is available from the BLAST web page, the standalone BLAST executables, or via the network BLAST client (see below).

2) Standalone BLAST executables: The Standalone BLAST executables are command line programs which run BLAST searches against local downloaded copies of the NCBI BLAST databases. The programs will handle either a single large file with multiple FASTA query sequences, or you can create a script to send multiple files one at a time. The executables are available for a wide variety of platforms, including many "flavors" of UNIX (LINUX, Solaris, etc.) Windows PC and even Mac OSX.

The Standalone executables are available at the anonymous FTP location:
<ftp://ftp.ncbi.nih.gov/blast/executables/> There is information on the Standalone BLAST executables available in the README file at
<ftp://ftp.ncbi.nih.gov/blast/documents/blast.txt> which is also bundled with the downloaded binaries.

3) BLAST Network Client 'blastcl3': The BLAST 2.0 Network client will allow you to submit a single file of FASTA sequences over an internet connection to the NCBI BLAST databases. You submit searches through the client to the NCBI servers and do not need to download the database locally. The BLAST Network client executables are located at:
<ftp://ftp.ncbi.nlm.nih.gov/blast/executables/> There are blastcl3 executables for various UNIX platforms, PC Windows and Macintosh.

Q: How can I write a program to submit jobs to NCBI's BLAST servers?

By using the URLAPI. Documentation also available in postscript and PDF.

Q: How can I limit my BLAST search based on Organism?

The option to limit a search to organism and even taxonomic classification is part of the "Limit by Entrez Search" option on most standard BLAST search pages. There is a pull down menu to select the most common organisms found in GenBank and also a field to input the species name, or classification (example: "eubacteria"). Using this option will cause your query sequences to be compared only to sequences in our databases from that organism.

There are also several "specialized" BLAST Pages devoted to different organisms on the main BLAST web page.

How can I limit my search to a subset of database sequences?

You can use the "Limit by Entrez Search" option found on most Standard BLAST search pages to run an Entrez search and have your query sequence compared to the results of this search. For example, if you wanted to limit your search to all phosphorylase sequences from mouse you could enter the following valid Entrez search strategy in the Limit by Entrez field of the BLAST search page: phosphorylase AND "Mus musculus"[Organism]

Q: Is it possible to search for a motif or pattern with BLAST?

There are two general approaches to this type of questions. First do you wish to find if motifs exist in your query sequence, or do you have a known motif and wish to find other proteins or nucleotides with this motif?

In the first case, finding motifs in your query sequence can be done for proteins using the CDD (Conserved Domain Database) and CDART (Conserved Domain Architecture Retrieval Tool) tools. CDD allows you to compare your protein to a database of alignments and

profiles representing protein domains conserved in molecular evolution as well as 3-dimensional protein structures in the MMDB database. These tools use popular protein motif databases, PFam (<http://pfam.wustl.edu/>) and Smart (<http://smart.embl-heidelberg.de>) in addition to the MMDB database.

For conditions of the second case if you have a known motif and wish to identify other proteins with this motif you can use PHI-BLAST. PHI-BLAST searches take a motif pattern and protein sequence as input and then compares these to the NCBI protein databases looking for other proteins which contain conserved regions similar to the motif entered.

For nucleotides it is only possible to search with short query sequences representing your motif or region of interest with the Nucleotide BLAST "Search for short nearly exact matches" service from the main BLAST web page. This can find other sequences which contain similar nucleotide patterns. However there are no database of nucleotide patterns which can identify patterns in your nucleotide query sequence.

You may also be interested in checking out other molecular biology web sites, such as those mentioned in the Other Molecular Biology Resources section at the end of this FAQ, for motif searching software.

Q: How do I perform a similarity search with a short peptide/nucleotide sequence?

There is a special page with pre-set parameters for searching with short sequences. You can access this page by clicking the "Search for short nearly exact matches" link on the main BLAST web page.

Essentially for these searches, the Expect value has been increased and the word size decreased to optimise for short hits which generally score a large E value require smaller word sizes to initiate formation of the HSP for extension. In addition, for proteins, the matrix "PAM30" becomes the default which optimises hits to smaller sequences which have a lower percentage of evolutionary drift in general.

Q: Can I use BLAST to compare to two or more sequences in a multiple sequence alignment?

✓ You can use the BLAST 2 Sequences service to compare two nucleotide or two protein sequences against each other using the Gapped BLAST algorithm. This will allow you to perform a BLAST search between the two sequences allowing for the introduction of gaps (deletions and insertions) in the resulting alignment. Remember that BLAST is a "local" alignment program and does not make global alignments between sequences to calculate total percent homologies.

To compare one sequence against a specific sequence or set of sequences, you can also use a separate multiple sequence alignment program. There are many such software tools available to do this. You may also be interested in checking out other molecular biology web sites, such as those mentioned in the Other Molecular Biology Resources section at the end of this FAQ.

Q: What is the Expect (E) value?

The Expect value (E) is a parameter that describes the number of hits one can "expect" to

see just by chance when searching a database of a particular size. It decreases exponentially with the Score (S) that is assigned to a match between two sequences. Essentially, the E value describes the random background noise that exists for matches between sequences. For example, an E value of 1 assigned to a hit can be interpreted as meaning that in a database of the current size one might expect to see 1 match with a similar score simply by chance. This means that the lower the E-value, or the closer it is to "0" the more "significant" the match is. However, keep in mind that searches with short sequences, can be virtually identical and have relatively high EValue. This is because the calculation of the E-value also takes into account the length of the Query sequence. This is because shorter sequences have a high probability of occurring in the database purely by chance. For more details please see the calculations in the BLAST Course.

The Expect value can also be used as a convenient way to create a significance threshold for reporting results. You can change the Expect value threshold on most main BLAST search pages. When the Expect value is increased from the default value of 10, a larger list with more low-scoring hits can be reported.

Q: What is low-complexity sequence?

Regions with low-complexity sequence have an unusual composition and this can create problems in sequence similarity searching (Wootton & Federhen, 1996). Low-complexity sequence can often be recognized by visual inspection. For example, the protein sequence PPCDPPPPPKDKKKKDDGPP has low complexity and so does the nucleotide sequence AAATAAAAAAATAAAAAAT. Filters are used to remove low-complexity sequence because it can cause artifactual hits (please also see Q: After running a search why do I see a string of "X"s (or "N"s) in my query sequence that I did not put there?)

In BLAST searches performed without a filter, often certain hits will be reported with high scores only because of the presence of a low-complexity region. Most often, this type of match cannot be thought of as the result of homology shared by the sequences. Rather, it is as if the low-complexity region is "sticky" and is pulling out many sequences that are not truly related.

Other Molecular Biology Resources:

The on-line BLAST Course was written by Dr. Stephen Altschul and discusses the basics of the Gapped BLAST algorithm. In addition the full text of the 1997 Nucleic Acids Research paper "Gapped BLAST and PSI-BLAST: a new generation of protein database search programs" is also available on-line.

Other links:

European Bioinformatics Institute (EBI) BioCatalog
Indiana University IUBio Archive
Sequence manipulation site

Troubleshooting

Q: Causes for "No significant similarity found".

Below are several reasons that a BLAST search can result in the "No significant similarity found" message.

Short Sequences: There is a special BLAST optimized for searching with small sequences.

Go to the main [BLAST web page](#) and select the "Search for short nearly exact matches" link for Nucleotide - Nucleotide or Protein Protein sections.

Filtering: BLAST filters regions of low-complexity (for a description of low-complexity see "[What is low-complexity sequence?](#)" below). If your sequence contains large regions of "low complexity" it may not significant hits to the database. You can turn off filtering by setting the "Filter" option to "None" using the pull down tab.

Query Format: Another reason you may see the "No Significant Similarity found" message is using the wrong type of sequence in your search.

1) Accession/GI Number or FASTA. Check that you have the Input Data set to the correct format for your Query. Set the pull down menu to "Accession number or Gi" to search with GenBank accession numbers or Gi numbers. Set to FASTA for raw amino acid or nucleotide sequences. For more information on FASTA format, [click here](#).

2) Sequence type and Program combination. You can search with an amino acid query sequence using the blastp and tblastn programs. With nucleotide query sequences you can use blastn, blastx, and tblastx. Please note that tblastx program cannot be used with the nr database on the BLAST Web page.

For more information on the BLAST programs, [click here](#).

Q: Why does my search timeout on the BLAST servers?

Certain combinations of BLAST searches with large sequences against large databases can cause the BLAST servers to timeout. This has to do with a limit on the server CPU's which prevents sequences which generate many HSPs from hoarding server resources.

However there are some things you can do to prevent timeout and generate results from large sequences.

- Some sequences contain large regions of ALU repeats. In this case you can select the "Human Repeat" filtering option on the main BLAST search page. This will mask repeat regions which generate a large number of biologically uninteresting hits to the databases.

- Increase the Word Size to 20 - 25. With a default Word Size of 7, the BLAST algorithm finds initial HSPs of 7 bases in length and begins extension of these from either end. In a large sequence this can generate 100's of initial HSPs between the query sequence and even a single large genomic sequence in the databases. Increasing the Word Size to 25 makes the initial HSP smaller, limiting the number small initial fragments to be extended.

- Decrease the Expect value to 1.0 or lower. Many hits from large sequences are to many small fragments in the database. The expect value for these searches is such that decreasing the expect value will eliminate these results, and concentrate on results which are more likely to contain large coding regions and genomic fragments.

If you are still seeing a "timeout" error message after making the above changes, please contact blast-help@ncbi.nlm.nih.gov with the RID of your search.

Q: Why do I get the message "ERROR:BLASTSetUpSearch: Unable to calculate Karlin-Altschul params, check querysequence" ?

This will happen if your entire query sequence has been masked by low complexity filtering. You will need to turn filtering off to get hits. For further information on filtering, please read the sections of the BLAST FAQs on Q: What is low-complexity sequence? and also Q: After running a search why do I see a string of "X"s (or "N"s) in my query sequence that I did not put there?

Q: Why do I get the message "ERROR: Blast: No valid letters to be indexed"?

You may have accidentally entered an accession number in the search box without changing the input selection from "Sequence in FASTA format" to "Accession or gi". You will also see this error message if too many ambiguity codes (R,Y,K,W,N, etc. for nucleotides) are present in your query sequence. Although BLAST allows ambiguity codes, be aware that these will always contribute a negative score in nucleic acid searches. Thus, sequences such as degenerate PCR primers with ambiguity codes may not find any significant hits even though they may be designed from sequences that are present in the database.

Q: After running a search why do I see a string of "X"s (or "N"s) in my query sequence that I did not put there?

You are seeing the result of automatic filtering of your query for low-complexity sequence that is performed to prevent artifactual hits. The filter substitutes any low-complexity sequence that it finds with the letter "N" in nucleotide sequence (e.g., "NNNNNNNNNNNNNN") or the letter "X" in protein sequences (e.g., "XXXXXXXXXX"). Low-complexity regions can result in high scores that reflect compositional bias rather than significant position-by-position alignment (Wootton and Federhen, 1996). Filter programs can eliminate these potentially confounding matches from the blast reports, leaving regions whose BLAST statistics reflect the specificity of their pairwise alignment. Queries searched with the blastn program are filtered with DUST. The other BLAST programs use SEG.

Q: How can I see low-similarity matches when there are many strong hits to my query sequence? Often, when the query is a member of a large sequence family, the summary hit list and the alignments returned only contain very high scoring hits. To look at low-similarity matches, you must increase the maximum number of results returned. On the BLAST Web pages, often it is sufficient to increase the size of the summary hit list and the number of alignments shown using the menus on the Advanced pages. However, it is possible to increase the lists even further using the Other Advanced Options box on the Advanced BLAST pages. For BLAST 2.0, "-v 2000", for example, will increase the number of descriptions returned in the summary hit list to 2000. The option "-b 2000" will similarly increase the number of alignments returned.

Q: I have heard that I will be penalized if I send a large number of sequences to the servers?

The NCBI WWW BLAST server is a shared resource and it would be unfair for a few users to monopolize it. To prevent this, the server keeps track of how many queries are in the queue for each user and penalizes those users with many queries in the queue. This is done by calculating a 'Time of Execution' (TOE). If a user has only one query in the queue, then the TOE is set to the current time. As a user adds more queries to the queue, then the TOE is set to the current time, plus 60 seconds for every query in the queue. An example would be if a user sent in five requests one after the other without waiting for any to be worked on, then the TOE's for the requests would be:

1st request: current time

2nd request: current time + 60 seconds
3rd request: current time + 120 seconds
4th request: current time + 180 seconds
5th request: current time + 240 seconds

The BLAST server works through requests in the order of earliest to latest TOE. A query will be executed before it's TOE, if there are no other queries with an earlier TOE. Users with large numbers of queries are encouraged to use the BLAST servers at off-peaks hours, which are from 8 p.m. to 8 a.m. (EST).

[Disclaimer](#)
[Privacy statement](#)
[Accessibility](#)
Valid [XHTML 1.0](#), [CSS](#).



More Information

glossary

BLAST
information


Query
tutorial

BLAST
tutorial

BLAST
guide

PSI-BLAST
tutorial

Glossary



[back to Education](#)

Alignment

The process of lining up two or more sequences to achieve maximal levels of identity (and conservation, in the case of amino acid sequences) for the purpose of assessing the degree of similarity and the possibility of homology.

Algorithm

A fixed procedure embodied in a computer program.

Bioinformatics

The merger of biotechnology and information technology with the goal of revealing new insights and principles in biology.

Bit score

The value S' is derived from the raw alignment score S in which the statistical properties of the scoring system used have been taken into account. Because bit scores have been normalized with respect to the scoring system, they can be used to compare alignment scores from different searches.

BLAST

Basic Local Alignment Search Tool. (Altschul et al.) A sequence comparison algorithm optimized for speed used to search sequence databases for optimal local alignments to a query. The initial search is done for a word of length "W" that scores at least "T" when compared to the query using a substitution matrix. Word hits are then extended in either direction in an attempt to generate an alignment with a score exceeding the threshold of "S". The "T" parameter dictates the speed and sensitivity of the search. For additional details, see one of the BLAST tutorials (Query or BLAST) or the narrative guide to BLAST.

BLOSUM

Blocks Substitution Matrix. A substitution matrix in which scores for each position are derived from observations of the frequencies of substitutions in blocks of local alignments in related proteins. Each matrix is tailored to a particular evolutionary distance. In the BLOSUM62 matrix, for example, the alignment from which scores were derived was created using sequences sharing no more than 62% identity. Sequences more identical than 62% are represented by a single sequence in the alignment so as to avoid over-weighting closely related family members. (Henikoff and Henikoff)

Conservation

Changes at a specific position of an amino acid or (less commonly, DNA) sequence that preserve the physico-chemical properties of the original residue.

Domain

A discrete portion of a protein assumed to fold independently of

BLAST
information
tutorial
guide

PSI-BLAST
tutorial

More Information
similarity searching
rules of thumb
glossary
reference list

the rest of the protein and possessing its own function.

DUST

A program for filtering low complexity regions from nucleic acid sequences.

E value

Expectation value. The number of different alignments with scores equivalent to or better than S that are expected to occur in a database search by chance. The lower the E value, the more significant the score.

FASTA

The first widely used algorithm for database similarity searching. The program looks for optimal local alignments by scanning the sequence for small matches called "words". Initially, the scores of segments in which there are multiple word hits are calculated ("init1"). Later the scores of several segments may be summed to generate an "initn" score. An optimized alignment that includes gaps is shown in the output as "opt". The sensitivity and speed of the search are inversely related and controlled by the "k-tup" variable which specifies the size of a "word". (Pearson and Lipman)

Filtering

Also known as Masking. The process of hiding regions of (nucleic acid or amino acid) sequence having characteristics that frequently lead to spurious high scores. See SEG and DUST.

gap

A space introduced into an alignment to compensate for insertions and deletions in one sequence relative to another. To prevent the accumulation of too many gaps in an alignment, introduction of a gap causes the deduction of a fixed amount (the gap score) from the alignment score. Extension of the gap to encompass additional nucleotides or amino acid is also penalized in the scoring of an alignment.

Global Alignment

The alignment of two nucleic acid or protein sequences over their entire length.

H

H is the relative entropy of the target and background residue frequencies. (Karlin and Altschul, 1990). H can be thought of as a measure of the average information (in bits) available per position that distinguishes an alignment from chance. At high values of H, short alignments can be distinguished by chance, whereas at lower H values, a longer alignment may be necessary. (Altschul, 1991)

Homology

Similarity attributed to descent from a common ancestor.

HSP

High-scoring segment pair. Local alignments with no gaps that achieve one of the top alignment scores in a given search.

Identity

The extent to which two (nucleotide or amino acid) sequences are invariant.

K

A statistical parameter used in calculating BLAST scores that can be thought of as a natural scale for search space size. The value K is used in converting a raw score (S) to a bit score (S').

lambda

A statistical parameter used in calculating BLAST scores that can be thought of as a natural scale for scoring system. The value lambda is used in converting a raw score (S) to a bit score (S').

Local Alignment

The alignment of some portion of two nucleic acid or protein sequences

Low Complexity Region (LCR)

Regions of biased composition including homopolymeric runs, short-period repeats, and more subtle overrepresentation of one or a few residues. The SEG program is used to mask or filter LCRs in amino acid queries. The DUST program is used to mask or filter LCRs in nucleic acid queries.

Masking

Also known as Filtering. The removal of repeated or low complexity regions from a sequence in order to improve the sensitivity of sequence similarity searches performed with that sequence.

Motif

A short conserved region in a protein sequence. Motifs are frequently highly conserved parts of domains.

Multiple Sequence Alignment

An alignment of three or more sequences with gaps inserted in the sequences such that residues with common structural positions and/or ancestral residues are aligned in the same column. Clustal W is one of the most widely used multiple sequence alignment programs

Optimal Alignment

An alignment of two sequences with the highest possible score.

Orthologous

Homologous sequences in different species that arose from a common ancestral gene during speciation; may or may not be responsible for a similar function.

P value

The probability of an alignment occurring with the score in question or better. The p value is calculated by relating the observed alignment score, S, to the expected distribution of HSP scores from comparisons of random sequences of the same length and composition as the query to the database. The most highly significant P values will

be those close to 0. P values and E values are different ways of representing the significance of the alignment.

PAM

Percent Accepted Mutation. A unit introduced by Dayhoff et al. to quantify the amount of evolutionary change in a protein sequence. 1.0 PAM unit, is the amount of evolution which will change, on average, 1% of amino acids in a protein sequence. A PAM(x) substitution matrix is a look-up table in which scores for each amino acid substitution have been calculated based on the frequency of that substitution in closely related proteins that have experienced a certain amount (x) of evolutionary divergence.

Paralogous

Homologous sequences within a single species that arose by gene duplication.

Profile

A table that lists the frequencies of each amino acid in each position of protein sequence. Frequencies are calculated from multiple alignments of sequences containing a domain of interest. See also PSSM.

Proteomics

Systematic analysis of protein expression of normal and diseased tissues that involves the separation, identification and characterization of all of the proteins in an organism.

PSI-BLAST

Position-Specific Iterative BLAST. An iterative search using the BLAST algorithm. A profile is built after the initial search, which is then used in subsequent searches. The process may be repeated, if desired with new sequences found in each cycle used to refine the profile. Details can be found in this discussion of PSI-BLAST. (Altschul et al.)

PSSM

Position-specific scoring matrix; see profile. The PSSM gives the log-odds score for finding a particular matching amino acid in a target sequence.

Query

The input sequence (or other type of search term) with which all of the entries in a database are to be compared.

Raw Score

The score of an alignment, S , calculated as the sum of substitution and gap scores. Substitution scores are given by a look-up table (see PAM, BLOSUM). Gap scores are typically calculated as the sum of G , the gap opening penalty and L , the gap extension penalty. For a gap of length n , the gap cost would be $G+Ln$. The choice of gap costs, G and L is empirical, but it is customary to choose a high value for G (10-15) and a low value for L (1-2).

Similarity

RAFFINOSE SYNTHASE GENES AND THEIR USE

BACKGROUND OF THE INVENTION

FIELD OF INVENTION

The present invention relates to raffinose synthase genes and their use.

DISCLOSURE OF THE RELATED ART

Raffinose family oligosaccharides are derivatives of sucrose, which are represented by the general formula:
 $\text{o-}\alpha\text{-D-galactopyranosyl-(1}\rightarrow\text{6)}_n\text{-o-}\alpha\text{-D-glucopyranosyl-(1}\rightarrow\text{2)-}\beta\text{-D-fluctofuranoside}$, and they are called "raffinose" when n is 1, "stachyose" when n is 2, "verbascose" when n is 3, and "ajugose" when n is 4.

It has been known that raffinose family oligosaccharides have an effect of giving good conditions of enterobacterial flora, if present at an appropriate amount in food. Therefore, raffinose family oligosaccharides have already been used as a functional food material for addition to some kinds of food and utilized in the field of specific health food. On the other hand, raffinose family oligosaccharides are neither digested nor absorbed in mammals such as human, but are assimilated and decomposed by enterobacteria to generate gases and to cause meteorism and absorption disorder. Therefore, it has been desired to appropriately regulate the amount of raffinose family oligosaccharides in food and feed.

Raffinose family oligosaccharides are synthesized



by the raffinose family oligosaccharide biosynthesis system beginning with sucrose in many plants. This biosynthesis system normally involves a reaction for the sequential addition of galactosyl groups from galactinol through an α (1 \rightarrow 6) bond to
5 the hydroxyl group attached to the carbon atom at 6-position of the D-glucose residue in a sucrose molecule. Raffinose synthase is the enzyme concerned in the reaction for producing raffinose by allowing a D-galactosyl group derived from galactinol to form the α (1 \rightarrow 6) bond with the hydroxyl group
10 attached to the carbon atom at 6-position of the D-glucose residue in a sucrose molecule in the first step of this biosynthesis system. It has been suggested that this enzyme constitutes a rate-limiting step in the above synthesis system, and therefore this enzyme is quite important in the control of biosynthesis
15 of raffinose family oligosaccharides.

Then, a method for controlling an expression level or activity of raffinose synthase in plants by utilizing a raffinose synthase gene is effective to control a biosynthesis system of raffinose family oligosaccharides in plants to increase
20 or decrease the production of raffinose in plants. Thus, a raffinose synthase gene which can be used in such a method has been desired.

OBJECTS OF THE INVENTION

25 The main object of the present invention is to provide novel raffinose synthase genes from plants.

This object as well as other objects and advantage of the present invention will become apparent to those skilled in the art from the following description.

5

~~SUMMARY OF THE INVENTION~~

~~Under these circumstances, the present inventors have intensively studied and succeeded in isolating novel genes encoding raffinose synthase from various plants. Thus, the present invention has been completed.~~

10

~~That is, the present invention provides:~~

~~1. A raffinose synthase gene which comprises a nucleotide sequence hybridizable with a nucleotide sequence selected from the group consisting of:~~

15

~~(a) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 1,~~

~~(b) the nucleotide sequence represented by SEQ ID NO: 2,~~

~~(c) a nucleotide sequence encoding the amino acid sequence of represented by SEQ ID NO: 3,~~

20

~~(d) the nucleotide sequence represented by the 236th to 2584th nucleotides in the nucleotide sequence represented by SEQ ID NO: 4,~~

~~(e) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 5,~~

25

~~(f) the nucleotide sequence represented by the 134th to 2467th nucleotides in the nucleotide sequence represented~~

by ~~SEQ ID NO: 6,~~

~~———— (g) a nucleotide sequence encoding the amino acid
sequence represented by SEQ ID NO: 7, and~~

~~———— (h) the nucleotide sequence represented by the 1st
5 to 1719th nucleotides in the nucleotide sequence represented
by SEQ ID NO: 8,~~

~~under stringent conditions, and encoding a protein being capable
of binding D-galactosyl group through α (1 \rightarrow 6) bond to the hydroxyl
group attached to the carbon atom at 6-position of the D-glucose
10 residue in a sucrose molecule to form raffinose.~~

~~———— 2. A raffinose synthase gene comprising a nucleotide
sequence encoding the amino acid sequence represented by SEQ
ID NO: 1.~~

~~———— 3. A raffinose synthase gene comprising the
15 nucleotide sequence represented by SEQ ID NO: 2.~~

~~———— 4. A raffinose synthase gene comprising a nucleotide
sequence encoding the amino acid sequence represented by SEQ
ID NO: 3.~~

~~———— 5. A raffinose synthase gene comprising the
20 nucleotide sequence represented by the 236th to 2584th
nucleotides in the nucleotide sequence represented by SEQ ID
NO: 4.~~

~~———— 6. A raffinose synthase gene comprising a nucleotide
sequence encoding the amino acid sequence represented by SEQ
25 ID NO: 5.~~

~~———— 7. A raffinose synthase gene comprising the~~

~~nucleotide sequence represented by the 134th to 2467th
nucleotides in the nucleotide sequence represented by SEQ ID
NO: 6.~~

5 ~~8. A raffinose synthase gene comprising a nucleotide
sequence encoding the amino acid sequence represented by SEQ
ID NO: 7.~~

~~9. A raffinose synthase gene comprising the
nucleotide sequence represented by the 1st to 1719th nucleotides
in the nucleotide sequence represented by SEQ ID NO: 8.~~

10 ~~10. A raffinose synthase gene comprising the
nucleotide sequence represented by SEQ ID NO: 4, SEQ ID NO:
6 or SEQ ID NO: 8.~~

15 ~~11. A nucleic acid comprising a partial nucleotide
sequence of the raffinose synthase gene of any one of the above
1 to 10.~~

~~12. A method for detecting a nucleic acid containing
a raffinose synthase gene which comprises detecting said nucleic
acid by hybridization using the labeled nucleic acid of the
above 11 as a probe.~~

20 ~~13. A method for amplifying a nucleic acid containing
a raffinose synthase gene which comprises amplifying said nucleic
acid by polymerase chain reaction (PCR) using the nucleic acid
of the above 11 as a primer.~~

25 ~~14. A method for obtaining a raffinose synthase gene
which comprises the steps of:~~

~~detecting a nucleic acid containing said raffinose~~

~~synthase gene by hybridization using the labeled nucleic acid of the above 11 as a probe, and~~

~~recovering the detected nucleic acid.~~

5 ~~15. A method for obtaining a raffinose synthase gene which comprises the steps of:~~

~~amplifying a nucleic acid containing said raffinose synthase gene by PCR using the nucleic acid of the above 11 as a primer, and~~

~~recovering the amplified nucleic acid.~~

10 ~~16. A nucleic acid comprising a nucleic acid containing the raffinose synthase gene of any one of the above 1 to 10 which is joined to a nucleic acid exhibiting promoter activity in a host cell.~~

15 ~~17. A vector comprising the raffinose synthase gene of any one of the above 1 to 10.~~

~~18. A transformant, wherein the raffinose synthase gene of any one of the above 1 to 10 is introduced into a host cell.~~

20 ~~19. A transformant, wherein the nucleic acid of the above 16 is introduced into a host cell.~~

~~20. A transformant, wherein the vector of the above 17 is introduced into a host cell.~~

~~21. The transformant of any one of the above 18 to 20, wherein the host is a microorganism.~~

25 ~~22. The transformant of any one of the above 18 to 20, wherein the host is a plant.~~

- ~~23. A method for producing a raffinose synthase which comprises the steps of:~~
- ~~culturing or growing the transformant of any one of the above 18 to 22 to produce the raffinose synthase, and~~
- 5 ~~collecting the raffinose synthase.~~
- ~~24. A raffinose synthase comprising the amino acid sequence represented by SEQ ID NO: 1.~~
- ~~25. A raffinose synthase comprising the amino acid sequence represented by SEQ ID NO: 3.~~
- 10 ~~26. A raffinose synthase comprising the amino acid sequence represented by SEQ ID NO: 5.~~
- ~~27. A raffinose synthase comprising the amino acid sequence represented by SEQ ID NO: 7.~~
- ~~The term "nucleic acid" used herein means an oligomer~~
- 15 ~~compound or a high molecular compound which is generally called "DNA" or "RNA".~~

DETAILED DESCRIPTION OF THE INVENTION

The gene engineering techniques described below can

20 be carried out, for example, according to methods described in "Molecular Cloning: A Laboratory Manual 2nd edition" (1989), Cold Spring Harbor Laboratory Press, ISBN 0-87969-309-6; "Current Protocols In Molecular Biology" (1987), John Wiley & Sons, Inc. ISBN 0-471-50338-X; "Current Protocols In Protein

25 Science" (1995), John Wiley & Sons, Inc. ISBN 0-471-11184-8.

The genes of the present invention can be obtained

from soybean, plants belonging to the families Chenopdiaceae such as beet, etc. and Cruciferae such as mustard, rapeseed, etc. Specific examples of the genes of the present invention include those comprising a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 1, the nucleotide sequence represented by SEQ ID NO: 2, a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 3, the nucleotide sequence represented by SEQ ID NO: 4 or by the 236th to 2584th nucleotides in the nucleotide sequence represented by SEQ ID NO: 4, a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 5, the nucleotide sequence represented by SEQ ID NO: 6 or by the 134th to 2467th nucleotides in the nucleotide sequence represented by SEQ ID NO: 6, a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 7, the nucleotide sequence represented by SEQ ID NO: 8 or by the 1st to 1719th nucleotides in the nucleotide sequence represented by SEQ ID NO: 8, and the like.

The genes of the present invention can be obtained, for example, by the following method.

That is, the genes of the present invention derived from soybean can be obtained, for example, by the following method.

For example, the gene can be obtained by a hybridization method using a nucleic acid having the nucleotide sequence represented by SEQ ID NO: 2 or its partial nucleotide sequence as a probe to detect a nucleic acid fragment which hybridizes

to the probe in DNAs derived from soybean, followed by isolating the detected nucleic acid.

In this method, first, a nucleic acid to be used as the probe is prepared. As such a nucleic acid, for example,
5 there is a nucleic acid composed of an oligonucleotide chemically synthesized by a conventional method on the basis of the nucleotide sequence of SEQ ID NO: 2. Specific example thereof includes a nucleic acid having the 800th to the 899th nucleotides in the nucleotide sequence represented by SEQ ID NO: 2.

10 Alternatively, the gene of the present invention derived from soybean can be obtained by the following method.

For example, tissue of soybean (*Glycine max*) is frozen in liquid nitrogen and ground physically with a mortar or other means into finely divided tissue debris powder. From the tissue
15 debris powder, RNA is extracted by a conventional method. A commercially available RNA extraction kit can be utilized in the extraction. RNA is recovered from thus-obtained RNA extract by ethanol precipitation. Poly-A tailed RNA is fractionated from thus-recovered RNA by a conventional method. A commercially
20 available oligo-dT column can be utilized in this fractionation.

cDNA is synthesized from the poly-A tailed RNA thus obtained by a conventional method. The synthesis can be carried out by using a commercially available cDNA synthesis kit. DNA is amplified by PCR using the above-obtained cDNA as the template
25 and primers designed and synthesized on the basis of the nucleotide sequence of SEQ ID NO: 2. More specifically, as the primers,

for example, there are primers 11 (SEQ ID NO: 9) and 12 (SEQ ID NO: 10) shown in List 1 hereinafter. When PCR is carried out by using these primers and as the template cDNA derived from soybean, the genes of the present invention derived from soybean, e.g., the "raffinose synthase gene having a nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 1" and the "raffinose synthase gene having the nucleotide sequence of SEQ ID No: 2" can be obtained.

The amplified DNA can be cloned according to a conventional method, for example, described in "Molecular Cloning: A Laboratory Manual 2nd edition" (1989), Cold Spring Harbor Laboratory Press; or "Current Protocols In Molecular Biology" (1987), John Wiley & Sons, Inc. ISBN 0-471-50338-X.

Alternatively, cloning can be carried out, for example, by using a commercially available cloning kit such as TA cloning kit (Invitrogen) and a commercially available plasmid vector such as pBluescript II (Stratagene). The nucleotide sequence of the DNA clone can be determined by dideoxy terminating method such as that described by F. Sanger, S. Nicklen, A.R. Coulson, Proceedings of National Academy of Science U.S.A. (1977), 74, pp. 5463-5467. For example, preferably, a commercially available kit such as ABI PRISM Dye Terminator Cycle Sequencing Ready Reaction Kit manufactured by Perkin-Elmer can be used.

List 1

Primer 11 (SEQ ID NO: 9): ccaatctgat catgcttgtg ccgaa
25mer

Primer 12 (SEQ ID NO: 10): ggaacaaagt tatgcactat
tatttaaggt 30mer

5 The genes of the present invention derived from a
Chenopdiaceae plant such as beet can be obtained by the following
method.

For example, tissue of a Chenopdiaceae plant such
as beet (*Beta vulgaris*) is frozen in liquid nitrogen and ground
physically with a mortar or other means into finely divided
tissue debris powder. From this tissue debris powder, RNA is
10 extracted by a conventional method. A commercially available
RNA extraction kit can be utilized in the extraction. RNA is
recovered from the thus-obtained RNA extract by ethanol
precipitation. From the recovered RNA, poly-A tailed RNA is
fractionated by a conventional method. A commercially available
15 oligo-dT column can be utilized in this fractionation. cDNA
is synthesized from the poly-A tailed RNA thus obtained by a
conventional method. The synthesis can be carried out by
utilizing a commercially available cDNA synthesis kit. DNA
is amplified by PCR using the above-obtained cDNA as the template
20 and primers designed and chemically synthesized on the basis
of the nucleotide sequence of SEQ ID NO: 4. More specifically,
as the primers, for example, there are primers 21 (SEQ ID NO:
11) and 22 (SEQ ID NO: 12) shown in List 2 hereinafter. When
PCR is carried out by using these primers and as the template
25 cDNA derived from beet, the genes of the present invention derived
from beet, e.g., the "raffinose synthase gene having a nucleotide

sequence encoding the amino acid sequence of SEQ ID NO: 3,"
and the "raffinose synthase gene having a nucleotide sequence
of SEQ ID No: 4" can be obtained. According to a particular
purpose, the PCR primers can also be designed and synthesized
5 on the basis of the nucleotide sequence of SEQ ID NO: 4. For
example, in order to amplify the "raffinose synthase gene having
the nucleotide sequence represented by the 236th to the 2584th
nucleotides in the nucleotide sequence represented by SEQ ID
NO: 4", preferably, oligonucleotides having the nucleotide
10 sequences represented by primers 23 (SEQ ID NO: 13) and 24 (SEQ
ID NO: 14) in List 2 below are synthesized and used as the primers.

The amplified DNA can be cloned according to a
conventional method, for example, described in "Molecular
Cloning: A Laboratory Manual 2nd edition" (1989), Cold Spring
15 Harbor Laboratory Press; or "Current Protocols In Molecular
Biology" (1987), John Wiley & Sons, Inc. ISBN 0-471-50338-X.

Alternatively, cloning can be carried out by using a commercially
available cloning kit such as TA cloning kit (Invitrogen) and
a commercially available plasmid vector such as pBluescript
20 II (Stratagene). The nucleotide sequence of the DNA clone can
be determined, for example, by dideoxy terminating method such
as that described by F. Sanger, S. Nicklen, A.R. Coulson,
Proceedings of National Academy of Science U.S.A. (1977), 74,
pp. 5463-5467. For example, preferably, a commercially
25 available kit such as ABI PRISM Dye Terminator Cycle Sequencing
Ready Reaction Kit manufactured by Perkin-Elmer can be used.

List 2

Primer 21 (SEQ ID NO: 11): ctaccaaatt ccacaactta aagttca |
27mer

Primer 22 (SEQ ID NO: 12): ggaataataa gcttcacaca |
5 tactgtactc tc 32mer

Primer 23 (SEQ ID NO: 13): atgggtccaa gctttagcaa |
ggaaaattcc 30mer

Primer 24 (SEQ ID NO: 14): tcaaaataag tactcaacag |
tggtaaaacc 30mer

10 The genes of the present invention derived from
Cruciferae plants such as mustard (*Brassica juncea*) and rapeseed
(*Brassica napus*) can be obtained by the following method.

For example, tissue of a Cruciferae plant such as
mustard or rapeseed is frozen in liquid nitrogen and ground
15 physically with a mortar or other means into finely divided
tissue debris powder. From the tissue debris powder, RNA is
extracted by a conventional method. A commercially available
RNA extraction kit can be utilized in the extraction. The RNA
is recovered from thus-obtained RNA extract by ethanol
20 precipitation. Poly-A tailed RNA is fractionated from the RNA
thus recovered by a conventional method. A commercially
available oligo-dT column can be utilized in the fractionation.

cDNA is synthesized from the poly-A tailed RNA thus obtained
by a conventional method. The synthesis can be carried out
25 by using a commercially available cDNA synthesis kit. DNA are
amplified by PCR using the above-obtained cDNA as a template

and primers designed and chemically synthesized on the basis of the nucleotide sequence of SEQ ID NO: 6. For example, when PCR is carried out by using cDNA derived from mustard (*Brassica juncea*) as the template and primers 33 (SEQ ID NO: 17) and 34 (SEQ ID NO: 18) shown in List 3 hereinafter, the genes from Cruciferae plants of the present invention, e.g., the "raffinose synthase gene having a nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 5," and the "raffinose synthase gene having the nucleotide sequence represented by the 1st to 2654th nucleotides in the nucleotide sequence represented by SEQ ID NO: 6" can be obtained. According to a particular purpose, the PCR primers can also be designed and synthesized on the basis of the nucleotide sequence of SEQ ID NO: 6. For example, in order to amplify DNA encoding the open reading frame region of the "raffinose synthase gene having a nucleotide sequence encoding a protein having the amino acid sequence of SEQ ID NO: 5", and the "raffinose synthase gene having the nucleotide sequence represented by the 134th to the 2467th nucleotides of SEQ ID NO: 6", preferably, oligonucleotides having the nucleotide sequences represented by primers 35 (SEQ ID NO: 19) and 36 (SEQ ID NO: 20) in List 3 are synthesized and used as the primers.

The amplified DNA can be cloned according to a conventional method, for example, described in "Molecular Cloning: A Laboratory Manual 2nd edition" (1989), Cold Spring Harbor Laboratory Press; or "Current Protocols In Molecular

Biology" (1987), John Wiley & Sons, Inc. ISBN 0-471-50338-X.

Alternatively, cloning can be carried out, for example, by using a commercially available TA cloning kit (Invitrogen) or a commercially available plasmid vector such as pBluescript II (Stratagene). The nucleotide sequence of the DNA clone can be determined by dideoxy terminating method such as that described by F. Sanger, S. Nicklen, A.R. Coulson, Proceedings of National Academy of Science U.S.A. (1977), 74, pp. 5463-5467. For example, preferably, the commercially available ABI PRISM Dye Terminator Cycle Sequencing Ready Reaction Kit of Perkin-Elmer can be used.

List 3

Primer 31 (SEQ ID NO: 15):	ttggaagaga	agacgccgcc	
gggaatcgtc 30mer			
Primer 32 (SEQ ID NO: 16):	ttaagccccg	gcgagagctc	
tggccggaca 30mer			
Primer 33 (SEQ ID NO: 17):	accaatccaa	aatctcatca	
aataatcgca 30mer			
Primer 34 (SEQ ID NO: 18):	aaataatagg	ggcagtacaa	
attacaccac 30mer			
Primer 35 (SEQ ID NO: 19):	atggctccac cgagcgtaat taaatccga		
29mer			
Primer 36 (SEQ ID NO: 20):	ctaaaactca	tacttaatag	
aagacaaacc 30mer			

Then, a nucleic acid having a partial nucleotide sequence of the gene of the present invention (hereinafter referred to as "the gene fragment") which is obtained by the

above-described method is labeled and then used as a probe in a hybridization method. The probe can be hybridized to, for example, DNA derived from soybean, a Chenopdiaceae plant or a Cruciferae plant to detect a nucleic acid having the probe specifically bound thereto, thereby detecting a nucleic acid having the raffinose synthase gene.

As the DNA derived from soybean, a Chenopdiaceae plant such as beet or a Cruciferae plant such as mustard or rapeseed, for example, a cDNA library or a genomic DNA library of these plants can be used. The gene library may also be a commercially available gene library as such or a library constructed according to a conventional library construction method, for example, described in "Molecular Cloning: A Laboratory Manual 2nd edition" (1989), Cold Spring Harbor Laboratory Press; "Current Protocols In Molecular Biology" (1987), John Wiley & Sons, Inc. ISBN 0-471-50338-X.

As the hybridization method, for example, plaque hybridization or colony hybridization can be employed, and they are selected depending upon the kind of vector used in the construction of a library. More specifically, when the library to be used is constructed with a phage vector, a suitable host microorganism is mixed with the phage of the library under infectious conditions to obtain transformants. The transformant is further mixed with a soft agar medium, and the mixture is plated on an agar medium. Thereafter, the mixture is cultured at 37°C until a plaque of an appropriate size appears.

When the library to be used is constructed with a plasmid vector, the plasmid is introduced into a suitable host microorganism to form transformants. The transformant obtained is diluted to a suitable concentration and the dilution is plated on an agar medium, after which it is cultured at 37°C until a colony of an appropriate size appears. In either case of the above libraries, a membrane filter is placed on the surface of the agar medium after the above cultivation, so that the phage or transformant is transferred to the membrane. This membrane is denatured with an alkali, followed by neutralization, and for example, when a nylon membrane is used, the membrane is irradiated with ultraviolet light, so that DNA of the phage or transformant is fixed on the membrane. This membrane is then subjected to a hybridization method wherein the gene fragment which has a partial nucleotide sequence of the gene of the present invention and labeled by a conventional method (hereinafter referred to as "the labeled gene fragment") is used as a probe. For this method, reference may be made, for example, to D.M. Glover ed., "DNA cloning, a practical approach" IRL PRESS (1985), ISBN 0-947946-18-7. There are various reagents and temperature conditions to be used in the hybridization. For example, in general, prehybridization is carried out by immersion of the membrane in a prehybridization solution [6 x SSC (0.9 M NaCl, 0.09 M citric acid), 0.1 to 1 (w/v)% SDS, 100 µg/ml denatured salmon sperm DNA] and incubation at 65°C for 1 hour. Then, hybridization is carried out by addition and mixing of the labeled

gene fragment thereto and incubating the membrane at 42 to 68°C for 4 to 16 hours.

In the present invention, the "stringent conditions" are those wherein incubation is carried out, for example, at
5 65 to 68°C in the above hybridization.

After hybridization, the membrane is taken out and is washed with 2 x SSC containing 0.1 to 1 (w/v)% SDS, further rinsed with 0.2 x SSC containing 0.1 to 1 (w/v)% SDS, and then dried. The membrane is analyzed, for example, by autoradiography
10 or other techniques to detect the position of the probe on the membrane, thereby detecting the position on the membrane of a nucleic acid having a nucleotide sequence homologous to that of the probe used. The clone corresponding to the position of the nucleic acid thus detected on the membrane is identified
15 on the original agar medium and the positive clone is selected so that the clone having the nucleic acid can be isolated. The same procedures of detection are repeated to purify the clone having the nucleic acid.

Alternatively, a commercially available kit such as
20 GENE TRAPPER cDNA Positive Selection System kit (GibcoBRL) can be used. In this method, first, a single-stranded DNA library is hybridized with the biotinylated gene fragment (i.e., probe), followed by adding streptoavidin-bound magnet beads and mixing. From the mixture, the streptoavidin-bound magnetic beads are
25 collected with a magnet, so that single-stranded DNA having a nucleotide sequence homologous to that of the probe used,

which has been bound to these beads through the gene fragment, biotin and streptoavidin, is collected and detected. The single-stranded DNA collected can be converted into a double-strand form by reaction with a suitable DNA polymerase using a suitable oligonucleotide as a primer.

As described above, a nucleic acid containing raffinose synthase gene can be obtained by detecting a nucleic acid hybridizable to the gene fragment in DNAs of a gene library derived from soybean, a Chenopodiaceae plant or a Cruciferae plant, purifying a clone having the nucleic acid and isolating phage or plasmid DNA from the clone. By preparing the restriction map or determining the nucleotide sequence of the nucleic acid thus obtained according to a conventional method, the nucleic acid containing the gene of the present invention can be confirmed.

For example, the gene of the present invention from a Chenopodiaceae plant can be confirmed by the following point:

The amino acid encoded by the nucleotide sequence thus determined has 75% or more homology to the amino acid sequence represented by the 103rd to 208th amino acids in the amino acid sequence of SEQ ID NO: 3;

80% or more homology to the amino acid sequence represented by the 255th to 271st amino acids in the amino acid sequence of SEQ ID NO: 3;

70% or more homology to the amino acid sequence represented by the 289th to 326th amino acids in the amino acid sequence of SEQ ID NO: 3; or

70% or more homology to the amino acid sequence represented by the 610th to 696th amino acids in the amino acid sequence of SEQ ID NO: 3.

5 The gene of the present invention from a Cruciferae plant can be confirmed, for example, by the following point:

The amino acid sequence encoded by the nucleotide sequence determined has 75% or more homology to the amino acid sequence represented by the 111th to 213th amino acids in the amino acid sequence of SEQ ID NO: 5;

10 80% or more homology to the amino acid sequence represented by the 260th to 275th amino acids in the amino acid sequence of SEQ ID NO: 5;

70% or more homology to the amino acid sequence represented by the 293rd to 325th amino acids in the amino acid sequence of SEQ ID NO: 5; or

15

70% or more homology to the amino acid sequence represented by the 609th to 695th amino acids in the amino acid sequence of SEQ ID NO: 5.

The "homology" used herein means the proportion of the number of amino acids in a region, which are identical to those in a different region to be compared, to the number of the entire amino acids in the former region, upon comparing regions having similarity in two amino acid sequences. In this respect, it is preferred that the region having similarity contains more amino acids. Such homology of amino acid sequences can be evaluated by using a commercially available gene analysis

20

25

software such as GENETIX (Software Kaihatu K.K.).

Further, according to the same manner as described above, a nucleic acid containing raffinose synthase gene can be detected by hybridization to DNA from the desired organism using the gene fragment as a probe to detect a nucleic acid to which the probe specifically binds (hereinafter referred to as the detection method of the present invention). The gene fragment used herein can be chemically synthesized according to a conventional method on the basis of the nucleotide sequence represented by SEQ ID NO: 2, 4, 6 or 8. Alternatively, it can be prepared by PCR using as primers oligonucleotides chemically synthesized according to a conventional method on the basis of the nucleotide sequence represented by SEQ ID NO: 2, 4, 6 or 8.

The gene fragment may be a part of the non-translated region of the raffinose synthase gene as well as the open reading frame thereof. For example, an oligonucleotide having the same nucleotide sequence as a part of that of 5'-upstream side such as the 1st to 235th nucleotides in the nucleotide sequence of SEQ ID NO: 4, the 1st to 133rd nucleotides in the nucleotide sequence of SEQ ID NO: 6 and the like, or a part of that of 3'-downstream side such as the 2588th to 2675th nucleotides in the nucleotide sequence of SEQ ID NO: 4, the 2468th to 2676th nucleotides in the nucleotide sequence of SEQ ID NO: 6 and the like.

When PCR is carried out by using the gene fragment

as primers, it is possible to amplify a nucleic acid containing raffinose synthase gene from DNA derived from the desired organism (hereinafter referred to as the amplification method of the present invention).

5 More specifically, for example, oligonucleotides having the nucleotide sequences of the gene fragment are designed and chemically synthesized according to a conventional method.

In general, it is preferred that the number of nucleotides is more from a viewpoint that the specificity of annealing is
10 ensured. It is, however, also preferred that the number of nucleotides is not so many from viewpoints that the primers themselves are liable to have a higher structure giving possible deterioration of the annealing efficiency and that complicated procedures are required in the purification after the synthesis.

15 Normally, oligonucleotides composed of 15 to 50 bases are preferred. In this respect, based on the codon table showing the correspondence of amino acids encoded by codons, a mixture of primers can also be synthesized by using a mixture of plural bases so that a residue at a specified position in a primer
20 is changed to different bases according to the variation of codons which can encode one certain amino acid. Alternatively, for example, a base such as inosine which can form a base pair with plural bases can be used instead of the above mixture of plural bases.

25 Coding Table

Phe: UUU, UUC Ser: UCU, UCC, UCA, UCG, AGU, AGC

Tyr: UAU, UAC Cys: UGU, UGC
 Stop: UAA, UAG, UGA Trp: UGG
 Leu: UUA, UUG, CUU, CUC, CUA, CUG Pro: CCU, CCC, CCA,
 CCG His: CAU, CAC Gln: CAA, CAG Arg: CGU,
 5 CGC, CGA, CGG, AGA, AGG
 Ile: AUU, AUC, AUA Thr: ACU, ACC, ACA, ACG
 Asn: AAU, AAC Lys: AAA, AAG
 Met: AUG
 Val: GUU, GUC, GUA, GUG Ala: GCU, GCC, GCA, GCG
 10 Asp: GAU, GAC Gly: GGU, GGC, GGA, GGG
 Glu: GAA, GAG

In the above codon table, each codon is shown as the nucleotide sequence in mRNA and its right hand is the 5'-terminus.

U represents uracil base in RNA and corresponds to thymine base in DNA.

An oligonucleotide having the same nucleotide sequence as the coding strand of the double-stranded DNA of the gene of the present invention is called a "sense primer" and that having a nucleotide sequence complementary to the coding strand is called an "antisense primer".

A sense primer having the same nucleotide sequence as that of 5'-upstream side in the coding strand of the gene of the present invention, and an antisense primer having a nucleotide sequence complementary to the nucleotide sequence on the 3'-downstream side in this coding strand are used in combination for PCR reaction, for example, with a gene library,

genomic DNA or cDNA as the template to amplify DNA. As the gene library to be used, for example, there are a cDNA library and a genomic library derived from soybean, a Chenopdiaceae plant such as beet or a Cruciferae plant such as mustard or rapeseed, etc. The gene library may also be a library constructed according to a conventional library construction method, for example, described in "Molecular Cloning: A Laboratory Manual 2nd edition" (1989), Cold Spring Harbor Laboratory Press; "Current Protocols In Molecular Biology" (1987), John Wiley & Sons, Inc. ISBN 0-471-50338-X, or a commercially available gene library as such. As the genomic DNA or cDNA, for example, there are those prepared from soybean, a Chenopdiaceae plant such as beet or a Cruciferae plant such as mustard or rapeseed, etc. For example, PCR is carried out by using the primers 31 (SEQ ID NO: 15) and 32 (SEQ ID NO: 16) in the above List 3 and as the template cDNA derived from mustard to amplify DNA having the nucleotide sequence represented by the 749th to 1215th nucleotides in the nucleotide sequence of SEQ ID NO: 6. Further, PCR is carried out by using the primers and as the template cDNA derived from rapeseed to amplify DNA having the nucleotide sequence represented by the 1st to 467th nucleotides in the nucleotide sequence of SEQ ID NO: 8. The nucleic acid thus amplified can be confirmed by conventional electrophoresis. The nucleic acid can be cloned according a conventional method such as that described in "Molecular Cloning: A Laboratory Manual 2nd edition" (1989), Cold Spring Harbor laboratory Press or

"Current Protocols in Molecular Biology" (1987), John Wiley & Sons, Inc., ISBN 0-471-50338-X. For the nucleic acid, its restriction map is prepared or its nucleotide sequence is determined by a conventional method, so that the nucleic acid containing raffinose synthase gene or a part thereof can be identified. When the nucleic acid contains a part of raffinose synthase, PCR can be carried out on the basis of its nucleotide sequence to amplify the nucleic acid containing the 5'-upstream side nucleotide sequence or the 3'-downstream side nucleotide sequence. That is, based on the nucleotide sequence of the above-obtained nucleic acid, an antisense primer is designed and synthesized for amplification of the 5'-upstream side part, and a sense primer is designed and synthesized for amplification of the 3'-downstream side part. The nucleotide sequence of the 5'-upstream side part or 3'-downstream side part of the nucleotide sequence already obtained can be determined by the RACE method using these primers and a commercially available kit such as Marathon Kit of Clontech. The full length raffinose synthase gene can be obtained by synthesizing new primers based on both terminal sequences in the nucleotide sequence thus determined and carrying out PCR again.

The above detection method of the present invention can also be used in the analysis of genotypes of a plant such as soybean, a Chenopodiaceae plant or a Cruciferae plant, etc. More specifically, for example, a genomic DNA derived from soybean, a Chenopodiaceae plant or a Cruciferae plant is prepared

according to a conventional method, for example, described in "Cloning and Sequence (Plant Biotechnology Experiment Manual)" complied under the supervision of Itaru Watanabe, edited by Masahiro Sugiura, published by Noson Bunka-sha, Tokyo (1989).

5 The genomic DNA is digested with at least several kinds of restriction enzymes, followed by electrophoresis. The electrophoresed DNA is blotted on a filter according to a conventional method. This filter is subjected to hybridization with a probe prepared from DNA having the gene fragment by a
10 conventional method, and DNA to which the probe hybridizes is detected. The DNAs detected are compared in length between different varieties of a specified plant species. The differences in length make possible the analysis of differences in phenotypic characteristics accompanied with the expression
15 of raffinose family oligosaccharides between these varieties. Furthermore, when the DNAs detected by the above method are compared in length between the gene recombinant plant and the non-gene recombinant plant of the same variety, the former plant can be distinguished from the latter plant by the detection
20 of hybridizing bands greater in number or higher in concentration for the former plant than for the latter plant. This method can be carried out according to the RFLP (restriction fragment length polymorphism) method, for example, described in "Plant PCR Experiment Protocols" complied under the supervision of
25 Ko Shimamoto and Takuji Sasaki, published by Shujun-sha, Tokyo (1995), ISBN 4-87962-144-7, pp. 90-94.

Further, the amplification method of the present invention can be used for an analysis of genes of soybean, a Chenopdiaceae plant or a Cruciferae plant, etc. More specifically, for example, the amplification method of the present invention is carried out by using plant genomic DNA prepared from soybean, a Chenopdiaceae plant or a Cruciferae plant to amplify DNA. The amplified DNA is mixed with a formaldehyde solution, followed by heat denaturing at 85°C for 5 minutes and then quickly cooling on ice. A sample thereof is subjected to electrophoresis on, for example, 6 (w/v)% polyacrylamide gel containing 0 (v/v)% or 10 (v/v)% of glycerol. For this electrophoresis, a commercially available electrophoresis apparatus such as that for SSCP (Single Strand Conformation Polymorphism) can be used and the electrophoresis can be carried out with maintaining the gel at a constant temperature, for example, at 5°C, 25°C, 37°C, etc. From the electrophoresed gel, DNA is detected, for example, by a method such as silver staining method with a commercially available reagent. From the differences of behavior between the varieties in the electrophoresis of the DNA detected, a mutation in the raffinose synthase gene is detected, and an analysis is carried out for differences caused by the mutation in phenotypic characteristics accompanied with the expression of raffinose family oligosaccharides. This method can be carried out according to the SSCP method, for example, described in "Plant PCR Experiment Protocols" compiled under the supervision of

Ko Shimamoto and Takuji Sasaki, published by Shujun-sha, Tokyo (1995), ISBN 4-87962-144-7, pp. 141-146.

The analysis of the plant gene from soybean, a Chenopdiaceae plant or a Cruciferae plant by the above detection
5 method or amplification method of the present invention can be used not only for the analysis of differences in phenotypic characteristics accompanied with the expression of raffinose family oligosaccharides, but also, for example, for the selection of clones having the desired characters upon production of a
10 novel variety of soybean, a Chenopdiaceae plant or a Cruciferae plant. Further, it can also be used for identification of a clone thus produced and having the characters derived from a recombinant plant upon producing a plant variety using the recombinant plant.

15 For expression of the gene of the present invention in cells of a host, preferably, a nucleic acid comprising a nucleic acid fragment which contains the gene of the present invention, and a nucleic acid fragment which has a promoter activity in the host cells and joined to the former nucleic
20 acid fragment (hereinafter referred to as the expression nucleic acid of the present invention) can be used.

The nucleic acid fragment having promoter activity in the expression nucleic acid of the present invention is not limited to a specific one, so long as it is functionable in
25 a host to be transformed. For example, there are synthetic promoters functionable in *Escherichia coli*, such as *E. coli*

lactose operon promoter, *E. coli* tryptophan operon promoter and tac promoter, etc.; yeast alcohol dehydrogenase gene (ADH) promoter, adenovirus major late (Ad.ML) promoter, SV40 early promoter, baculovirus promoter and the like. When the host
5 is a plant, the promoter may include, for example, T-DNA derived constitutive promoters such as nopaline synthase gene (NOS) promoter, octopine synthase gene (OCS) promoter, etc.; plant virus-derived promoters such as cauliflower mosaic virus (CaMV)-derived 19S and 35S promoters; inducible promoters such
10 as phenylalanine ammonia-lyase (PAL) gene promoter, chalcone synthase (CHS) gene promoter, pathogenesis-related protein (PR) gene promoter, etc. Furthermore, vector pSUM-GY1 (see JP-A 06-189777/1994) can also be used, which has a promoter giving specific expression in a specified plant tissue, e.g., a promoter
15 of soybean-derived seed storage protein glycinin gene (JP-A 6-189777).

Furthermore, a nucleic acid fragment having a terminator activity can be joined to the expression nucleic acid of the present invention. In this case, it is generally
20 preferred that the expression nucleic acid of the present invention is constructed so that the nucleic acid fragment having a terminator activity is positioned downstream the raffinose synthase gene. The terminator to be used is not particularly limited, so long as it is functionable in cells of a host to
25 be transformed. For example, when the host is a plant, there are T-DNA derived constitutive terminators such as nopaline

synthase gene (NOS) terminator, etc.; plant derived terminators such as terminators of allium virus GV1 or GV2, and the like.

The expression nucleic acid of the present invention can be introduced into a host cell according to a conventional gene engineering technique to obtain a transformant. If
5 necessary, the expression nucleic acid of the present invention can be inserted into a vector having a suitable marker depending upon a particular transformation technique for introduction of the nucleic acid into a host cell.

10 A vector into which the expression nucleic acid of the present invention is inserted can be introduced into a microorganism according to a conventional method, for example, described in "Molecular Cloning: A Laboratory Manual 2nd
15 edition" (1989), Cold Spring Harbor laboratory Press or "Current Protocols in Molecular Biology" (1987), John Wiley & Sons, Inc., ISBN 0-471-50338-X. The microorganism transformed with the vector can be selected on the basis of a selection marker such as antibiotic resistance, auxotrophy or the like. In case that
20 the gene of the present invention is joined to the downstream of an inducible promoter (e.g., tac promoter) in the translatable form in the selected microorganism (e.g., *E. coli* transformant), a translated product of the gene of the present invention can be expressed under conventional culture and inducible conditions and can be recovered as a peptide or a protein.

25 The raffinose synthase activity of the translated product of the gene of the present invention thus prepared can

be measured by, for example, a method described in L. Lehle and W. Tanner, Eur. J. Biochem., 38, 103-110 (1973) to identify the translated product having the "capability of binding D-galactosyl group through α (1 \rightarrow 6) bond to the hydroxyl group attached to the carbon atom at 6-position of the D-glucose residue in sucrose molecule". More specifically, for example, the gene of the present invention is cloned in pGEX-4T3 (Pharmacia) to obtain a plasmid containing the expression nucleic acid of the present invention. The resultant plasmid is introduced into, for example, *E. coli* HB101 strain to obtain a transformant. The resultant transformant is culture overnight and 1 ml of the culture is inoculated into 100 ml of LB culture medium. It is incubated at 37°C for about 3 hours and IPTG (isopropylthio- β -D-galactoside) is added at a final concentration of 1 mM, followed by further incubation for 5 hours. Cells are recovered from the culture broth by centrifugation and are suspended by addition of 10 times of the cell weight of 100 mM Tris-HCl (pH 7.4), 1 mM EDTA, 5 mM DTT (dithiothreitol), 1 mM PMSF (phenylmethanesulfonyl fluoride) and 1 mM benzamide. The suspension is sonicated with an ultrasonic disrupter (Branson) to disrupt the cells. The disrupted cell suspension is centrifuged to recover a soluble protein solution. The resultant protein solution is added to a reaction mixture containing at final concentrations of 100 mM Tris-HCl (pH 7.4), 5 mM DTT (dithiothreitol), 0.01% BSA, 200 μ M sucrose, 5 mM galactinol and 31.7 μ M [14 C] sucrose. The reaction mixture is

incubated at 37°C, followed by addition of 1.5 times in volume of ethanol and stirring. Insoluble materials are removed by centrifugation, the supernatant is spotted on, for example, a HPTLC cellulose thin layer chromatography plate (Merck, HPTLC plates cellulose) and then the plate is developed with
5 n-butanol-pyridine-water-acetic acid (60 : 40 : 30 : 3). The developed plate is dried and analyzed with an imaging analyzer (FUJIX Bio-Image Analyzer BAS-2000II manufactured by Fuji Film) to determine [¹⁴C] raffinose produced to measure the raffinose
10 synthase activity.

In addition, the translated product as prepared above can also be used as an antigen for producing an antibody. The antibody thus produced can be used for, for example, detection and determination of the gene of the present invention in a
15 crude protein extract prepared from an organism such as a plant.

When the host is a plant, the vector into which the gene of the present invention is inserted can be introduced into plant cells by a conventional means such as *Agrobacterium* infection method (JP-B 2-58917 and JP-A 60-70080),
20 electroporation into protoplasts (JP-A 60-251887 and JP-B 5-68575) or particle gun method (JP-A 5-508316 and JP-A 63-258525).

The plant cell transformed by the introduction of the vector can be selected on the basis of a selection marker, for example, resistance to an antibiotic such as kanamycin or hygromycin.

25 From the plant cell thus transformed, a transformed plant can be regenerated by a conventional plant cell cultivation method,

for example, described in "Plant Gene Manipulation Manual (How to Produce Transgenic Plants)" written by Uchimiya, 1990, Kodan-sha Scientific (ISBN 4-06-153513-7), pp. 27-55. Furthermore, the collection of seeds from the transformed plant
5 also makes it possible to proliferate the transformed plant. In addition, crossing between the transformed plant obtained and the non-transformed plant makes it possible to produce progenic plants with the characters of the transformed plant.

As gene engineering techniques in soybean, basically,
10 the above general techniques can be employed. More specifically, "transformation of soybean plant strain by particle gun" described in EP 301749, gene introduction methods. for example, described in Torisky, R.S., Kovacs, L., Avdiushko, S., Newman, J.D., Hunt, A.G. and Collins, G.B., "Development of a binary
15 vector system for plant transformation based on the supervirulent *Agrobacterium tumefaciens* strain Chry5", Plant Cell Rep., (1997), 17, p. 102-108, etc. can be employed.

As gene engineering techniques in a *Chenopodiaceae* plant, basically, the above general techniques can be employed.
20 More specifically, gene introduction methods, for example, described in M. Mannerlof, S. Tuveesson, P. Steen and P. Tenning, "Transgenic sugar beet tolerant to glyphosate", Euphytica (1997), 94, p 83-91, B.K. Konwar, "Agrobacterium tumefaciens-Mediated Genetic Transformation of Sugar Beet (*Beta vulgaris* L.)", J.
25 Plant Biochemistry & Biotechnology (1994), 3, p. 37-41 can be employed.

As gene engineering techniques in a Cruciferae plant, basically, the above general techniques can be employed. More specifically, the gene introduction can be carried out according to a method, for example, described in J. Fry, A. Barnason and
5 R.B. Horsch, "Transformation of Brassica napus with Agrobacterium tumefaciens based vectors", Plant Cell Reports (1987), 6, 321-325.

For example, when gene introduction is carried out by *Agrobacterium* infection method, first, the above-described
10 expression nucleic acid of the present invention is inserted into a binary vector. The resultant vector can be introduced into, for example, *Agrobacterium tumefaciens* LBA 4404 strain which has been converted into a competent state by treatment with calcium chloride. A transformant can be selected by an
15 appropriate selection method according to the selection marker gene of the vector, for example, cultivation of a strain containing the vector in a culture medium containing an antibiotic in case that the selection marker gene is that giving resistance to the antibiotic such as kanamycin. The resultant transformed
20 *Agrobacterium* strain can be culture in a liquid culture medium, for example, LB medium.

Soybean, a Chenopodiaceae plant or a Cruciferae plant can be transformed by using thus obtained *Agrobacterium* transformant culture broth as described below. For example,
25 seeds from soybean, beet, rapeseed or mustard is sowed aseptically in, for example, 1/2 MS medium containing 2% sucrose and 0.7%

agar. After about 1 week, cotyledons and petioles of the germinated plant are cut off with a scalpel aseptically and transplanted in, for example, MS medium containing 3% sucrose, 0.7% agar, 4.5 μ M BA, 0.05 μ M 2,4-D and 3.3 μ M AgNO₃ and cultured
5 for one day. The cotyledons and petioles thus precultured are transferred to 1000-fold dilution of the above *Agrobacterium* culture broth and allowed to stand for 5 minutes. The cotyledons and petioles are transferred to the same medium as that of the preculture again and cultured for about 3 to 4 days. The
10 cotyledones and petioles thus cultured are transferred to, for example, MS medium containing 3% sucrose, 4.5 μ M BA, 0.05 μ M 2,4-D, 3.3 μ M AgNO₃ and 500 mg/liter cefotaxim, followed by shaking for 1 day to remove microbial cells. The resultant cotyledons and petioles are transferred to, for example, MS
15 medium containing 3% sucrose, 0.7% agar, 4.5 μ M BA, 0.05 μ M 2,4-D, 3.3 μ M AgNO₃ 100 mg/liter cefotaxim and 20 mg/liter kanamycin, followed by culturing for 3 to 4 weeks. Then, the cotyledons and petioles are transferred to, for example, MS medium containing 3% sucrose, 0.7% agar, 4.5 μ M BA, 0.05 μ M
20 2,4-D, 100 mg/liter cefotaxim and 20 mg/liter kanamycin and cultured. Culture in this medium is continued with subculturing every 3 to 4 weeks. When a shoot are regenerated, it is subcultured in, for example, MS medium containing 3% sucrose, 0.7% agar and 20 mg/liter kanamycin for 3 to 4 weeks. When the plant
25 makes roots, it is transferred to vermiculite-peat moss (1 : 1) and acclimatized by culturing at 21 to 22°C under day and

night conditions of 12 hours: 12 hours = day time : night.
As the plant grows, it is transferred to appropriate cultivation
soil to culture the plant. A genomic DNA is extracted from
the leaf of the regenerated plant according to the above method
5 and PCR is carried out by using as primers having partial
nucleotide sequences of the expression nucleic acid of the present
invention to confirm the insertion of the gene of the present
invention into the plant.

As described hereinabove, by introducing the gene
10 of the present invention into a plant, for example, soybean,
a Chenopdiaceae plant or a Cruciferae plant, it is possible
to vary the expression level and activity of raffinose synthase
in the plant to control the amount of raffinose family
oligosaccharides in the plant. The gene of the present invention
15 is useful in techniques for varying the expression level and
activity of raffinose synthase in soybean, a Chenopdiaceae plant
or a Cruciferae plant on the basis of gene homology, for example,
techniques such as homologous recombination and antisense
technique, cosuppression and the like.

20 The following examples further illustrate the present
invention in detail but are not to be construed to limit the
scope of the present invention.

Example 1

Preparation of cDNA Derived from Soybean

25 About 2 g of immature seeds of soybean (*Glycine max*)
Williams82 were frozen in liquid nitrogen and then ground with

a mortar, to which 20 ml of Isogen (Nippon Gene) was added, and the mixture was further thoroughly ground. The ground material was transferred into a centrifugation tube, to which 4 ml of chloroform was added, and the mixture was stirred with
5 a vortex mixer and then centrifuged at 6,500 x g for 10 minutes at 4°C. The water layer was collected, to which 10 ml of isopropanol was added, and the mixture was stirred and then centrifuged at 6,500 x g for 10 minutes at 4°C. The resulting precipitate was washed with 10 ml of 70% ethanol and then dissolved in 1 ml of
10 elution buffer (10 mM Tris-HCl/pH 7.5, 1 mM EDTA, 0.1% SDS).

The solution was allowed to stand at 60°C for 10 minutes and then centrifuged at 10,000 x g for 1 minute to remove insoluble matter. To the resulting supernatant was added an equivalent volume of Oligotex-dT30 (Takara), and the mixture was stirred
15 and then allowed to stand at 65°C for 5 minutes. Further, the mixture was placed on ice and allowed to stand for 3 minutes, to which 200 µl of 5 M NaCl was added, and the mixture was mixed and then allowed to stand at 37°C for 10 minutes. The mixture was then centrifuged at 10,000 x g for 3 minutes at 4°C. The
20 precipitate was collected and then suspended in 1 ml of TE buffer, and the suspension was allowed to stand at 65°C for 5 minutes. Further, the suspension was placed on ice and then allowing to stand for 3 minutes, followed by centrifugation at 10,000 x g for 3 minutes at 4°C to remove precipitate.

25 To the resulting supernatant were added 100 µl of 3M

sodium acetate and 2 ml of ethanol to precipitate and collect RNA. The collected RNA was washed twice with 70% ethanol and then dissolved in 20 μ l of sterilized water, which was used for the subsequent cDNA synthesis. The amount of RNA obtained was
5 determined by the measurement of absorbance at 260 nm.

For the cDNA synthesis, First Strand Synthesis Kit for RT-PCR (Amersham) and cDNA Synthesis Kit (Takara) were used, and all operations were made according to the protocol attached to kits.

10 Example 2

Cloning of Raffinose Synthase Gene from Soybean cDNA

PCR was carried out by using the cDNA obtained from immature seeds of soybean (*Glycine max*) Williams82 in Example 1 as a template and the primers designed on the basis of the
15 amino acid sequence of SEQ ID No: 1, i.e., primers having nucleotide sequences shown in List 4 (SEQ ID NOS: 21 and 22) below to amplify a DNA fragment. The PCR was carried out with Gene Amp PCR Systems 2400 and DNA Thermal Cycler Model 480 of Perkin-Elmer using Advantage KlenTaq cDNA Kit of Clontech. The reaction was carried
20 out by repeating the cycle for maintaining at 94°C for 1 minute, at 50°C for 3 minutes and then at 72°C for 3 minutes 40 times to amplify the DNA fragment. The amplified DNA fragment was cloned with TA cloning kit (Invitrogen), followed by sequence reaction using ABI PRISM Dye Terminator Cycle Sequencing Ready Reaction
25 Kit of Perkin-Elmer and analysis of the nucleotide sequence with a 373S DNA sequencer of ABI. Based on this sequence, primers

having nucleotide sequences shown in List 5 below were synthesized.

The synthesis of cDNA was carried out with Marathon Kit of Clontech using mRNA obtained from leaves of soybean Williams82 in Example 1. The cDNA obtained was ligated to an adapter contained in the kit with ligase. These operations were carried out according to the protocol attached to the kit. By using the adapter-ligated cDNA thus prepared, PCR was carried out with the primers shown in List 5 (SEQ ID NO: 23) according to the same manner as the above. The nucleotide sequence in terminal region of the gene was analyzed according to the protocol attached to the Marathon Kit of Clontech. As a result, the nucleotide sequence of SEQ ID NO: 2 was determined.

List 4

4-5-F primer (SEQ ID NO: 21):
15 cgatggatgg giaaittiat icaiccigai tgggaiatgt t 41mer
4-6-RV primer (SEQ ID NO: 22):
ggccacatit tiacia(ag)icc iatiggigci aa 32mer

List 5

5-SC-2 (SEQ ID NO: 23):
20 tggtactagg cgaaacaaga gtagctctga 30mer

Example 3

Preparation of cDNA derived from Chenopdiaceae Plant

About 2 g of leaves of beet (*Beta vulgaris*: haming) was frozen in liquid nitrogen and then ground with a mortar, to which 20 ml of Isogen (Nippon Gene) was added, and the mixture was further thoroughly ground. The ground material was

transferred into a centrifugation tube, to which 4 ml of chloroform was added, and the mixture was stirred with a vortex mixer and then centrifuged at 6,500 x g for 10 minutes at 4°C. The water layer was collected, to which 10 ml of isopropanol was added, and the mixture was stirred and then centrifuged at 6,500 x g for 10 minutes at 4°C. The resulting precipitate was washed with 10 ml of 70% ethanol and then dissolved in 180 µl of DEPC-treated sterilized water. The solution was allowed to stand at 55°C for 5 minutes and 20 µl of 5 M NaCl was added thereto. The resulting solution was purified using BIOMAG mRNA PURIFICATION KIT (PerSeptive Biosystems: Catalog No. 8-MB4003K).

To the resulting mRNA solution were added 3M sodium acetate and ethanol, and RNA was precipitated and collected. The collected RNA was washed twice with 70% ethanol and then dissolved in 20 µl of sterilized water, which was used for the subsequent cDNA synthesis. The amount of RNA obtained was determined by the measurement of absorbance at 260 nm.

For the cDNA synthesis, SMART PCR cDNA Synthesis Kit (Clontech) was used, and all operations were made according to the protocol attached to the kit.

Example 4

Analysis of Nucleotide Sequence of Raffinose Synthase Gene from Chenopodiaceae Plant

Synthetic DNA primers having the nucleotide sequences shown in List 6 (SEQ ID NOS: 24-27) below were synthesized. The PCR method was carried out with Gene Amp PCR Systems 2400

and DNA Thermal Cycler Model 480 of Perkin-Elmer using Advantage KlenTaq cDNA Kit of Clontech. The PCR was carried out with the above primers (SEQ ID NOS: 24-27) and cDNA of beet obtained in the above Example 3 by repeating the cycle for maintaining at 94°C for 1 minute, at 50°C for 3 minutes and then at 72°C for 3 minutes 40 times. As a result, the combinations of primers 6-3-F (SEQ ID NO: 24) and 6-8-RV (SEQ ID NO: 25) and primers 6-10-F (SEQ ID NO: 26) and 6-6-RV (SEQ ID NO: 27) gave an amplification of about 0.3 kb and 0.6 kb bands, respectively.

The amplified DNA fragments were cloned with TA cloning kit (Invitrogen), followed by sequence reaction using ABI PRISM Dye Terminator Cycle Sequencing Ready Reaction Kit of Perkin-Elmer and analysis of nucleotide sequence with a 373S DNA sequencer of ABI. Based on the resulting nucleotide sequences, synthetic DNA primers having nucleotide sequences shown in List 7 (SEQ ID NOS: 28-35) below were prepared and PCR was carried out using cDNA obtained from beet in Example 3 in the same manner as above.

As a result, DNA having the nucleotide sequence of SEQ ID NO: 4 was finally obtained from cDNA of beet.

List 6

6-3-F (SEQ ID NO: 24):
cgaggigggt gicciccigg ittigtiati atigaigaig gitggca 47mer

6-8-RV (SEQ ID NO: 25):
at(t/c)tt(a/g)tcia cigcia(a/g)(a/g)tc (t/c)tccatigt 29mer

6-10-F (SEQ ID NO: 26):
ggiaacitait gg(c/t)ticaigg itgicaiatg gticaigt 38mer

6-6-RV (SEQ ID NO: 27):

ggccacatit tiacia(a/g)icc iatiggigci aa 32mer

List 7

7-Sb-1 (SEQ ID NO: 28):

5 atctatttgt catgacgatg atccga 26mer

7-Sb-2RV (SEQ ID NO: 29):

ggccctcatt cccatattgg gatgatcctc 30mer

7-Sb-3RV (SEQ ID NO: 30):

aagcatgcca aacatacaca tgctcaacag 30mer

10 7-Sb-4RV (SEQ ID NO: 31):

agacccgggg aaagctttgg ggttactact 30mer

7-Sb-5 (SEQ ID NO: 32):

tgatgggaa actttataca ccctgact 28mer

7-Sb-6 (SEQ ID NO: 33):

15 gacatgttcc catctacaca cccttggtg 28mer

7-Sb-7 (SEQ ID NO: 34):

ccaatttatg ttagtgatgt tgttggaag 30mer

7-Sb-8RV (SEQ ID NO: 35):

tcgactccca gggtagaatt gtcac 26mer

20 Example 5

Preparation of cDNA Derived from Cruciferae Plant

About 2 g of leaves of mustard (*Brassica juncea*) was frozen in liquid nitrogen and then ground with a mortar, to which 20 ml of Isogen (Nippon Gene) was added, and the mixture was further thoroughly ground. The ground material was transferred into a centrifugation tube, to which 4 ml of chloroform

was added, and the mixture was stirred with a vortex mixer and then centrifuged at 6,500 x g for 10 minutes at 4°C. The water layer was collected, to which 10 ml of isopropanol was added, and the mixture was stirred and then centrifuged at 6,500 x g
5 for 10 minutes at 4°C. The resulting precipitate was washed with 10 ml of 70% ethanol and then dissolved in 180 µl of DEPC-treated sterilized water. The solution was allowed to stand at 55°C for 5 minutes and to which 20 µl of 5 M NaCl was added. The resulting solution was purified using BIOMAG mRNA PURIFICATION
10 KIT (PerSeptive Biosystems: Catalog No. 8-MB4003K).

To the resulting mRNA solution were added 3M sodium acetate and ethanol, and RNA was precipitated and collected. The collected RNA was washed twice with 70% ethanol and then dissolved in 20 µl of sterilized water, which was used for the
15 subsequent cDNA synthesis. The amount of RNA obtained was determined by the measurement of absorbance at 260 nm.

For the cDNA synthesis, SMART PCR cDNA Synthesis Kit (Clontech) was used, and all operations were carried out according to the protocol attached to the kit.

20 In the same manner as described in the above, mRNA was purified from immature seeds of rapeseed Westar (*Brassica napus*) and cDNA was synthesized.

Example 6

Isolation and Nucleotide Sequence Analysis of Raffinose
25 Synthase Gene derived from Cruciferae Plant

DNA primers having the nucleotide sequences shown in

List 8 (SEQ ID NOS: 36 and 37) below were synthesized. PCR was carried out with Gene Amp PCR Systems 2400 and DNA Thermal Cycler Model 480 of Perkin-Elmer using Advantage KlenTaq cDNA Kit of Clontech. The PCR was carried out with the above primers and cDNA of mustard obtained in Example 5 by repeating the cycle for maintaining at 94°C for 1 minute, at 50°C for 3 minutes and then at 72°C for 3 minutes 40 times. The reaction products were analyzed by agarose gel electrophoresis. As a result, an amplification of about the 1.2 kb bands was detected. The amplified DNA fragment was cloned with TA cloning kit (Invitrogen), followed by sequence reaction using ABI PRISM Dye Terminator Cycle Sequencing Ready Reaction Kit of Perkin-Elmer and nucleotide sequence analysis with a 373S DNA sequencer of ABI. Based on the resulting nucleotide sequence, synthetic primers having the nucleotide sequences shown in List 9 (SEQ ID NOS: 38 and 39) below were prepared and PCR was carried out using cDNAs from mustard (*Brassica juncea*) and rapeseed Westar (*Brassica napus*) obtained in Example 5 according to the same manner as the above. As a result, the nucleotide sequence represented by the 749th to 1215th nucleotides of SEQ ID NO: 6 and by the 1st to 467th nucleotides of SEQ ID NO: 8 were finally determined for cDNA from mustard (*Brassica juncea*) and rapeseed Westar (*Brassica napus*), respectively.

List 8

25 8-#1 (SEQ ID NO: 36):

cgattiaaig titggtggac iacicaitgg gtigg

35mer

8-#10RV (SEQ ID NO: 37):

caitgiacca titgicaicc itgia(ag)ccai taigticc 38mer

List 9

9-primer-1 (SEQ ID NO: 38):

5 gttagggttc atatgaacac cttcaagctc 30mer

9-primer-2RV (SEQ ID NO: 39):

caacggcgag atcttgcac gtcaac 26mer

Example 7

Nucleotide Sequence Analysis of Raffinose Synthase

10 Full-Length Gene Derived from Cruciferae Plant

Based on the nucleotide sequences obtained in Example 6, DNA primers having the nucleotide sequences shown in List 10 (SEQ ID NOS: 40-46) below were synthesized. The cDNAs from mustard (*Brassica juncea*) and rapeseed Westar (*Brassica napus*) obtained in the same manner as described in Example 5 were ligated to adapters contained in Marathon Kit of Clontech. By using the adapter-ligated cDNAs thus prepared, PCR was carried out with primers shown in List 10. 10-B-2RV (SEQ ID NO: 40), 10-B-3RV (SEQ ID NO: 41) and 10-B-4RV (SEQ ID NO: 42) primers were used for nucleotide analysis of 5'-termini, and 10-B-1 (SEQ ID NO: 43), 10-B-8 (SEQ ID NO: 44), 10-B-7 (SEQ ID NO: 45) and 10-B-6 (SEQ ID NO: 46) primers were used for nucleotide analysis of 3'-termini. The nucleotide sequences were analyzed according to the protocol attached to the Marathon Kit of clontech. As a result, the nucleotide sequence of SEQ ID NO: 6 and SEQ ID NO: 8 were determined from mustard (*Brassica juncea*) and rapeseed

Westar (*Brassica napus*), respectively.

List 10

	10-B-2RV (<u>SEQ ID NO: 40</u>):	
	ggattcgaca caaaccgcca cgtcatcgtc	30mer
5	10-B-3RV (<u>SEQ ID NO: 41</u>):	
	ccacgtgcac caccggaact tatcgac	27mer
	10-B-4RV (<u>SEQ ID NO: 42</u>):	
	aacatcgata ccatcggagt catgtccaat	30mer
	10-B-1 (<u>SEQ ID NO: 43</u>):	
10	gttagggttc atatgaacac cttcaagctc	30mer
	10-B-8 (<u>SEQ ID NO: 44</u>):	
	tctacgtctg gcacgcgctt tgcggctac	29mer
	10-B-7 (<u>SEQ ID NO: 45</u>):	
	gttgacgtca tccacatatt ggagatgttg t	31mer
15	10-B-6 (<u>SEQ ID NO: 46</u>):	
	gttatcgcta gcatggagca ctgtaatga	29mer

Example 8

Construction of Expression Vectors in Plant for
Raffinose Synthase Gene Derived from Cruciferae Plant

20 Based on the nucleotide sequence of raffinose synthase
gene from mustard obtained in Example 7, DNA primers having
the nucleotide sequences shown in List 11 (SEQ ID NOS: 47 and
48) were prepared. By using cDNA of mustard, PCR was carried
out in the same manner as described in Example 6. The amplified
25 DNA fragment was digested with SacI. The DNA fragment thus
digested was ligated to the vector pBI121(-) previously digested

with SacI by using Ligation Kit (Takara). Plasmid pBI121 (Clontech) were digested with BamHI and SacI, and ligated to linkers shown in List 12 (SEQ ID NOS: 49 and 50) to prepare the vector pBI121(-). The vector thus obtained was analyzed
5 by a restriction map and PCR using primers having nucleotide sequences shown in List 13 (SEQ ID NOS: 51-53), and confirmed the direction of inserted raffinose synthase gene. The vector whose raffinose synthase gene from mustard was inserted in the expressible direction was designated BjRS-Sac(+)-121 and the
10 one whose raffinose synthase gene from mustard was inserted in the reverse direction was designated BjRS-Sac(-)-121.

List 11

11-SacI-BjN (SEQ ID NO: 47):
aacgagctca atccaaaatc tcatacaata atcgc 35mer
15 11-SacI-BjintRV (SEQ ID NO: 48):
acaatagttg agggcggaag agtag 25mer

List 12

12-BamSac-(+)linker (SEQ ID NO: 49):
gatcgagctc gtgtcggatc cagct 25mer
20 12-BamSac-(-)linker (SEQ ID NO: 50):
ggatccgaca cgagctc 17mer

List 13

13-35S-3 (SEQ ID NO: 51):
cctcctcgga ttccattgcc cagctatctg 30mer
25 13-B-2RV (SEQ ID NO: 52):
ggattcgaca caaaccgcca cgtcatcgtc 30mer

13-B-8 (SEQ ID NO: 53):

tctacgtctg gcacgcgctt tgcggctac

29mer

Example 9

Transformation with Raffinose Synthase Gene Derived
5 from Cruciferae Plant

The vectors BjRS-Sac(+)-121 and BjRS-Sac(-)-121 prepared in Example 8 were used for the transformation of mustard (*Brassica juncea*) by the *Agrobacterium* infection method.

Agrobacterium tumefaciens (strain LBA4404 having
10 rifampicin and streptomycin resistance) previously converted into a competent state by calcium chloride treatment was transformed independently with two plasmids BjRS-Sac(+)-121 and BjRS-Sac(-)-121 prepared in Example 8. The transformants were selected on LB medium containing 50 µg/ml rifampicin and
15 25 µg/ml kanamycin by utilizing the kanamycin resistant character conferred by the kanamycin resistant gene (neomycin phosphotransferase, NPTII) of the introduced plasmids.

The transformant *Agrobacterium* obtained (*Agrobacterium tumefaciens* strain LBA4404: rifampicin and
20 streptomycin resistant) was cultured on LB medium containing 50 µg/ml rifampicin and 25 µg/ml kanamycin at 28°C for a whole day and night, and the culture was used for the transformation of mustard by the method described below.

The seeds of mustard were aseptically sowed on 1/2
25 MS medium containing 2% sucrose and 0.7% agar. After one week, cotyledons and petioles of sprouting plants were cut out with

a scalpel, and transferred to MS medium containing 3% sucrose, 0.7% agar, 4.5 μ M BA, 0.05 μ M 2.4-D and 3.3 μ M AgNO₃, followed by preculture for 1 day. The precultured cotyledons and petioles were transferred in a 1000-fold dilution of the Agrobacterium culture broth and allowed to stand for 5 minutes. The cotyledons and petioles were transferred again to the same medium as used in the preculture, and cultured for 3 to 4 days. The cultured cotyledons and petioles were transferred to MS medium containing 3% sucrose, 4.5 μ M BA, 0.05 μ M 2.4-D, 3.3 μ M AgNO₃ and 500 mg/l cefotaxim, and shaken for 1 day to remove microbial cells. The cotyledons and petioles thus treated were transferred to MS medium containing 3% sucrose, 0.7% agar, 4.5 μ M BA, 0.05 μ M 2.4-D, 3.3 μ M AgNO₃, 100 mg/l cefotaxim and 20 mg/l kanamycin, and cultured for 3 to 4 weeks. The cotyledons and petioles were transferred to MS medium containing 3% sucrose, 0.7% agar, 4.5 μ M BA, 0.05 μ M 2.4-D, 100 mg/l cefotaxim and 20 mg/l kanamycin, and cultivated. The cultivation on this medium was continued with subculturing at intervals of 3 to 4 weeks. When shoots are began to regenerate, these shoots are subcultured on MS medium containing 3% sucrose, 0.7% agar and 20 mg/l kanamycin, and cultivated for 3 to 4 weeks. The rooting plants are transferred to vermiculite : peat moss = 1 : 1, and cultivated at 21°C to 22°C in a cycle of day/night = 12 hours : 12 hours. With the progress of plant body growth, the plants are grown with cultivation soil.

BRIEF DESCRIPTION OF SEQUENCES

SEQ ID NO: 1 shows an amino acid sequence of a raffinose synthase protein encoded by the raffinose synthase gene of the present invention.

5 SEQ ID NO: 2 shows a nucleotide sequence of the raffinose synthase gene of the present invention.

SEQ ID NO: 3 shows an amino acid sequence of a raffinose synthase protein encoded by the raffinose synthase gene obtained from beet.

10 SEQ ID NO: 4 shows a cDNA nucleotide sequence of the raffinose synthase gene obtained from beet.

SEQ ID NO: 5 shows an amino acid sequence of a raffinose synthase protein encoded by the raffinose synthase gene obtained from mustard.

15 SEQ ID NO: 6 shows a cDNA nucleotide sequence of the raffinose synthase gene obtained from mustard.

SEQ ID NO: 7 shows an amino acid sequence of a raffinose synthase protein encoded by the raffinose synthase gene obtained from rapeseed.

20 SEQ ID NO: 8 shows a cDNA nucleotide sequence of the raffinose synthase gene obtained from rapeseed.

List 1:

25 The nucleotide sequences shown in List 1 (SEQ ID NOS: 9 and 10) are examples of the typical primers used in the amplification of a DNA fragment having a raffinose synthase gene. All of these sequences are based on the nucleotide sequence

of SEQ ID NO: 2. Primer 11 (SEQ ID NO: 9) is a sense primer
and Primer 12 (SEQ ID NO: 10) is an antisense primer. Depending
upon the purpose, recognition sequences for suitable restriction
enzymes can be added to the 5'-termini of these nucleotide
5 sequences.

List 2:

The nucleotide sequences shown in List 2 (SEQ ID NOS:
11-14) are examples of the typical primers used in the amplification
of a cDNA of a raffinose synthase gene. Primer 21 (SEQ ID NO:
10 11) is a sense primer corresponding to the 5'-terminus of the
beet-derived raffinose synthase gene. Primer 22 (SEQ ID NO:
12) is an antisense primer corresponding to the 3'-terminus.

Depending upon the purpose, recognition sequences for suitable
restriction enzymes can be added to the 5'-termini of these
15 nucleotide sequences.

Primer 23 (SEQ ID NO: 13) is a sense primer corresponding
to the N-terminus of the open reading frame, and primer 24 (SEQ
ID NO: 14) is an antisense primer corresponding to the C-terminus.

List 3:

20 Among the nucleotide sequences shown in List 3 (SEQ
ID NOS: 15-20), primers 31 (SEQ ID NO: 15) and 32 (SEQ ID NO:
16) are typical primers used in the amplification of a DNA having
the partial nucleotide sequence of a raffinose synthase gene.

Primer 31 (SEQ ID NO: 15) is a sense primer used in the amplification
25 of a DNA having the partial nucleotide sequence of a raffinose
synthase gene from mustard and rapeseed and primer 32 (SEQ ID

NO: 16) is an antisense primer. Depending upon the purpose, recognition sequences for suitable restriction enzymes can be added to the 5'-termini of these nucleotide sequences.

Primers 33 (SEQ ID NO: 17) and 34 (SEQ ID NO: 18) are the typical primers used in the amplification of a cDNA of a raffinose synthase gene of mustard. Primers 33 (SEQ ID NO: 17) and 34 (SEQ ID NO: 18) are both based on the nucleotide sequence of raffinose synthase gene in the non-translated region. Primer 33 (SEQ ID NO: 17) is a sense primer corresponding to the 5'-terminal non-translated region of the mustard-derived raffinose synthase gene. Primer 34 (SEQ ID NO: 18) is an antisense primer corresponding to the 3'-terminal non-translated region.

Primers 35 (SEQ ID NO: 19) and 36 (SEQ ID NO: 20) are typical primers used in the amplification of an open reading frame coding for the amino acid sequence of a raffinose synthase protein in the cDNA of a raffinose synthase gene. Primer 35 (SEQ ID NO: 19) is a sense primer corresponding to the 5'-terminus of the above open reading frame. Primer 36 (SEQ ID NO: 20) is an antisense primer corresponding to the 3'-terminus. Depending upon the purpose, recognition sequences for suitable restriction enzymes can be added to the 5'-termini of these nucleotide sequences.

List 4:

The nucleotide sequences shown in List 4 (SEQ ID NOS: 21 and 22) are of the primers used in the cloning of a DNA fragment having the present raffinose synthase gene. The base represented

by the symbol "i" is inosine. The bases shown in parentheses mean that a mixture of those bases is used in the synthesis.

The symbol "RV" as used after the primer number means that the primer referred to by this symbol has an antisense sequence.

5 List 5:

The nucleotide sequence shown in List 5 (SEQ ID NO: 23) is of the primer used in the analysis of a nucleotide sequence of the present raffinose synthase gene. 5-SC-2 (SEQ ID NO: 23) is used in the analysis of the present nucleotide sequence in the 3'-terminal region.

List 6

The nucleotide sequences shown in List 6 (SEQ ID NOS: 24-27) are of the primers used in the analysis of the present raffinose synthase gene of beet. The base represented by the symbol "i" is inosine. The bases shown in parentheses mean that a mixture of those bases was used in the synthesis. The symbol "RV" as used after the primer number means that the primer referred to by this symbol has an antisense sequence.

List 7

The nucleotide sequences shown in List 7 (SEQ ID NOS: 28-35) are of the primers synthesized on the partial nucleotide sequences of the beet raffinose synthase gene. The symbol "RV" as used after the primer number means that the primer referred to by this symbol has an antisense sequence.

25 List 8:

The nucleotide sequences shown in List 8 (SEQ ID NOS:

36 and 37) are of the primers used in the analysis of the cDNA nucleotide sequence of a raffinose synthase gene of mustard.

The base represented by the symbol "i" is inosine. The bases shown in parentheses mean that a mixture of those bases. The
5 symbol "RV" as used after the primer number means that the primer referred to by this symbol has an antisense sequence.

List 9:

The nucleotide sequences shown in List 9 (SEQ ID NOS: 38 and 39) are of the primers synthesized on the partial nucleotide
10 sequences of the mustard raffinose synthase gene. The symbol "RV" as used after the primer number means that the primer referred to by this symbol has an antisense sequence.

List 10:

The nucleotide sequences shown in List 10 (SEQ ID NOS: 40-46) are of the primers used in the analysis of the nucleotide
15 sequences of raffinose synthase gene of mustard and rapeseed. The symbol "RV" as used after the primer number means that the primer referred to by this symbol has an antisense sequence.

List 11:

20 The nucleotide sequences shown in List 11 (SEQ ID NOS: 47 and 48) are of the primers used in the amplification of 5'-terminal region of a mustard raffinose synthase gene. 11-SacI-BjN (SEQ ID NO: 47) is a primer whose SacI restriction site is added to the nucleotide sequence represented by the 4th
25 to 29th nucleotides in SEQ ID NO: 6. 11-SacI-BjintRV (SEQ ID NO: 48) is an antisense primer having a nucleotide sequence

corresponding to the nucleotide sequence represented by the 1164th to 1188th nucleotides in SEQ ID NO: 6.

List 12:

The nucleotide sequences shown in List 12 (SEQ ID NOS: 49 and 50) are of the adapters added to a mustard cDNA. These synthetic DNA take a double-stranded form when mixed together because they are complementary strands. This adapter has cohesive ends of cleavage sites for the restriction enzymes BamHI and SacI on both termini, and contains the restriction sites for the restriction enzymes BamHI and SacI in the double-stranded region.

List 13:

The nucleotide sequences shown in List 13 (SEQ ID NOS: 51-53) are of the primers used in the confirmation of inserting direction of the mustard-derived raffinose synthase gene. 13-35S-3 (SEQ ID NO: 51) is a primer of sense to 35S promoter. 13-B-2RV (SEQ ID NO: 52) is an antisense primer having the nucleotide sequence represented by the 593rd to 622nd nucleotides of SEQ ID NO: 6, 13-B-8 (SEQ ID NO: 53) is a sense primer having the nucleotide sequence represented by the 1110th to 1138th nucleotides in SEQ ID NO: 6.

As described hereinabove, according to the present invention, it is possible to provide raffinose synthase genes which can be utilized in techniques for varying expression level and activity of raffinose synthase in plants.

SEQUENCE LISTING FREE TEXT

SEQ ID NO: 9 to SEQ ID NO: 20: Designed oligonucleotide primer to obtain raffinose synthase gene.

5 SEQ ID NO: 21 and SEQ ID NO: 22: Designed oligonucleotide primer to obtain raffinose synthase gene, n is i, r is a or g.

SEQ ID NO: 23: Designated oligonucleotide primer to obtain raffinose synthase gene.

10 SEQ ID NO: 24 to SEQ ID NO: 27: Designed oligonucleotide primer to obtain raffinose synthase gene, n is i, y is t or c, r is a or g.

SEQ ID NO: 28 to SEQ ID NO: 35: Designed oligonucleotide primer to obtain raffinose synthase gene.

15 SEQ ID NO: 36 and SEQ ID NO: 37: Designed oligonucleotide primer to obtain raffinose synthase gene, n is i, r is a or g.

SEQ ID NO: 38 to SEQ ID NO: 48: Designed oligonucleotide primer to obtain raffinose synthase gene.

20 SEQ ID NO: 49 and SEQ ID NO: 50: Designed oligonucleotide linker to obtain raffinose synthase gene.

SEQ ID NO: 51 to SEQ ID NO: 53: Designed oligonucleotide primer to confirm direction of the inserted raffinose synthase gene.



- 57 -

SEQUENCE LISTING

5 <110> Eijiro WATANABE et al.; Sumitomo Chemical Company Limited

 <120> Raffinose Synthase Genes and Their Use

 <150> JP 10/120550

10 <151> 1998-04-30

 <150> JP 10/120551

 <151> 1998-04-30

15 <150> JP 10/345590

 <151> 1998-12-04

 <150> JP 10/351246

 <151> 1998-12-10

20 <160> 53

 <210> 1

 <211> 265

25 <212> PRT

 <213> Glycine max

 <400> 1

30 Gln Ser Asp His Ala Cys Ala Glu Phe His Ala Ala Ser Arg Ala Ile

 5 10 15

 Ser Gly Gly Pro Ile Tyr Val Ser Asp Ser Val Gly Lys His Asn Phe

 20 25 30

 Lys Leu Leu Lys Lys Leu Val Leu Pro Asp Gly Ser Ile Leu Arg Cys

 35 40 45

35 Gln His Tyr Ala Leu Pro Thr Arg Asp Cys Leu Phe Val Asp Pro Leu

 50 55 60

 His Asp Gly Lys Thr Met Leu Lys Ile Trp Asn Leu Asn Lys Cys Ser

 65 70 75 80

 Gly Val Leu Gly Leu Phe Asn Cys Gln Gly Gly Gly Trp Cys Pro Val

40 85 90 95

	Thr Arg Arg Asn Lys Ser Ser Ser Asp Tyr Ser His Ser Val Thr Cys	
	100 105 110	
	Phe Ala Ser Pro Gln Asp Ile Glu Trp Gly Lys Gly Lys His Pro Val	
	115 120 125	
5	Cys Ile Lys Gly Val Asp Val Phe Ala Val Tyr Met Phe Lys Asp Asp	
	130 135 140	
	Lys Leu Lys Leu Leu Lys Tyr Thr Glu Ser Val Glu Val Ser Leu Glu	
	145 150 155 160	
	Pro Phe Ser Cys Glu Leu Leu Thr Val Ser Pro Val Val Ile Leu Pro	
10	165 170 175	
	Arg Lys Ser Ile Gln Phe Ala Pro Ile Gly Leu Val Asn Met Leu Asn	
	180 185 190	
	Ser Gly Gly Ser Ile Met Ser Leu Glu Phe Asp Gln Gln Glu Asn Leu	
	195 200 205	
15	Ala Arg Ile Gly Val Arg Gly His Gly Glu Met Arg Val Phe Ala Ser	
	210 215 220	
	Glu Lys Pro Glu Ser Val Lys Ile Asp Gly Glu Ser Val Glu Phe Asp	
	225 230 235 240	
	Tyr Val Asp Arg Thr Val Arg Leu Gln Val Ser Trp Pro Cys Ser Ser	
20	245 250 255	
	Arg Leu Ser Val Val Glu Tyr Leu Phe	
	260 265	
	<210> 2	
25	<211> 928	
	<212> DNA	
	<213> Glycine max	
	<220>	
30	<221> CDS	
	<222> (2)... (799)	
	<400> 2	
	c caa tct gat cat gct tgt gcc gaa ttc cac gct gct tct aga gcc	46
35	Gln Ser Asp His Ala Cys Ala Glu Phe His Ala Ala Ser Arg Ala	
	5 10 15	
	att tct ggt gga cca att tat gta agc gac tct gtt gga aaa cac aac	94
	Ile Ser Gly Gly Pro Ile Tyr Val Ser Asp Ser Val Gly Lys His Asn	
	20 25 30	
40	ttc aag ttg ctt aag aag ctt gtt cta cct gat ggc tcc att ttg cgg	142

	Phe Lys Leu Leu Lys Lys Leu Val Leu Pro Asp Gly Ser Ile Leu Arg	
	35 40 45	
	tgt caa cat tat gca ctt ccc acc cga gac tgc tta ttt gta gat cct	190
	Cys Gln His Tyr Ala Leu Pro Thr Arg Asp Cys Leu Phe Val Asp Pro	
5	50 55 60	
	tta cat gat ggg aaa aca atg ctc aaa att tgg aac ctc aat aaa tgt	238
	Leu His Asp Gly Lys Thr Met Leu Lys Ile Trp Asn Leu Asn Lys Cys	
	65 70 75	
10	tcc ggg gtt ttg ggt ctg ttc aat tgc caa gga gga ggt tgg tgc cct	286
	Ser Gly Val Leu Gly Leu Phe Asn Cys Gln Gly Gly Gly Trp Cys Pro	
	80 85 90 95	
	gtt act agg cga aac aag agt agc tct gac tat tca cac tcc gtg act	334
	Val Thr Arg Arg Asn Lys Ser Ser Ser Asp Tyr Ser His Ser Val Thr	
	100 105 110	
15	tgc ttt gca agt cct caa gac att gaa tgg ggc aaa ggg aag cac cca	382
	Cys Phe Ala Ser Pro Gln Asp Ile Glu Trp Gly Lys Gly Lys His Pro	
	115 120 125	
	gtt tgc atc aaa ggg gtg gac gta ttt gct gtg tac atg ttt aag gac	430
	Val Cys Ile Lys Gly Val Asp Val Phe Ala Val Tyr Met Phe Lys Asp	
20	130 135 140	
	gac aag ttg aag ctg ctg aag tac aca gag agt gta gaa gtt tct ctt	478
	Asp Lys Leu Lys Leu Leu Lys Tyr Thr Glu Ser Val Glu Val Ser Leu	
	145 150 155	
	gag cct ttt agt tgt gag ctt ttg acc gtt tct cca gtg gtg atc tta	526
25	Glu Pro Phe Ser Cys Glu Leu Leu Thr Val Ser Pro Val Val Ile Leu	
	160 165 170 175	
	ccc aga aaa tca atc caa ttt gcc cca att gga ttg gta aac atg ctc	574
	Pro Arg Lys Ser Ile Gln Phe Ala Pro Ile Gly Leu Val Asn Met Leu	
	180 185 190	
30	aac tct ggg ggc tct att atg tca ttg gaa ttt gat caa cag gaa aat	622
	Asn Ser Gly Gly Ser Ile Met Ser Leu Glu Phe Asp Gln Gln Glu Asn	
	195 200 205	
	ttg gcg agg att ggg gtg aga gga cat ggg gaa atg agg gta ttt gca	670
	Leu Ala Arg Ile Gly Val Arg Gly His Gly Glu Met Arg Val Phe Ala	
35	210 215 220	
	tca gag aag cca gag agt gtc aag att gat gga gaa tct gtg gaa ttt	718
	Ser Glu Lys Pro Glu Ser Val Lys Ile Asp Gly Glu Ser Val Glu Phe	
	225 230 235	
40	gat tat gtt gat aga acc gtg agg ctc caa gtc tcg tgg cct tgt tct	766
	Asp Tyr Val Asp Arg Thr Val Arg Leu Gln Val Ser Trp Pro Cys Ser	

```

240                245                250                255
tcg agg ttg tcc gta gtc gag tat ttg ttc tga atcatgattt ggtgtccgag 819
Ser Arg Leu Ser Val Val Glu Tyr Leu Phe
                260                265
5  agagccgtgt aatgttcaca taaactgact taagtgcatt aagcaaatcc accttaaata 879
   atagtgcata actttgttcc aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 928

<210> 3
<211> 783
10 <212> PRT
   <213> Beta vulgaris L.

<400> 3
Met Ala Pro Ser Phe Ser Lys Glu Asn Ser Lys Thr Cys Asp Glu Val
15      5      10      15
Ala Asn His Asp Asp Cys Asn Thr Cys Pro Ile Ile Ser Leu Glu Glu
      20      25      30
Ser Asn Phe Met Val Asn Gly His Val Ile Leu Ser Gln Val Pro Ser
      35      40      45
20 Asn Ile Thr Ala Ile Ser Lys Met Gly Phe Asp Gly Leu Phe Val Gly
      50      55      60
Phe Asp Ala Pro Glu Pro Lys Ala Arg His Val Val Ser Val Gly Gln
      65      70      75      80
Leu Lys Gly Ile Pro Phe Met Ser Ile Phe Arg Phe Lys Val Trp Trp
25      85      90      95
Thr Thr His Trp Thr Gly Ser Asn Gly Arg Asp Leu Glu His Glu Thr
      100     105     110
Gln Ile Leu Ile Leu Asp Lys Ser Asp Glu Gly Leu Gly Arg Pro Tyr
      115     120     125
30 Ile Val Ile Leu Pro Leu Ile Glu Gly Pro Phe Arg Ala Ser Leu Gln
      130     135     140
Pro Gly Ser Val Asp Asp Tyr Val Asp Ile Cys Val Glu Ser Gly Ser
145     150     155     160
Thr Lys Val Val Gly Asp Ser Phe Arg Ala Val Leu Tyr Ile Arg Ala
35     165     170     175
Gly Pro Asp Pro Phe Lys Leu Ile Lys Asp Thr Met Lys Glu Val Gln
      180     185     190
Ala His Leu Gly Thr Phe Lys Leu Leu Asp Asp Lys Thr Pro Pro Gly
      195     200     205
40 Ile Val Asp Lys Phe Gly Trp Cys Thr Trp Asp Ala Phe Tyr Leu Lys

```

	210		215		220	
	Val Glu Pro Tyr Gly	Val Trp Glu Gly Val	Lys Gly Leu Val Glu Asn			
	225	230	235	240		
	Gly Val Pro Pro Gly	Leu Val Leu Ile Asp	Asp Gly Trp Gln Ser Ile			
5	245	250	255			
	Cys His Asp Asp Asp	Pro Ile Thr Asp	Gln Glu Gly Ile Asn Arg Thr			
	260	265	270			
	Ser Ala Gly Glu Gln	Met Pro Cys Arg Leu	Ile Lys Tyr Glu Glu Asn			
	275	280	285			
10	Phe Lys Phe Arg Asp	Tyr Lys Ser Pro Asn	Ile Met Gly His Glu Asp			
	290	295	300			
	His Pro Asn Met Gly	Met Arg Ala Phe Val	Arg Asp Leu Lys Glu Glu			
	305	310	315	320		
	Phe Lys Thr Val Glu	His Val Tyr Val Trp	His Ala Phe Thr Gly Tyr			
15	325	330	335			
	Trp Gly Gly Val Arg	Pro Asn Val Pro Gly	Leu Pro Glu Ala Gln Val			
	340	345	350			
	Val Thr Pro Lys Leu	Ser Pro Gly Leu Glu	Met Thr Met Glu Asp Leu			
	355	360	365			
20	Ala Val Asp Lys Ile	Val Asn Asn Gly Ile	Gly Leu Val Gln Pro Asp			
	370	375	380			
	Lys Ala Gln Glu Leu	Tyr Glu Gly Leu His	Ser His Leu Glu Asn Cys			
	385	390	395	400		
	Gly Ile Asp Gly Val	Lys Val Asp Val Ile	His Leu Leu Glu Met Met			
25	405	410	415			
	Ala Glu Asp Tyr Gly	Gly Arg Val Glu Leu	Ala Lys Thr Tyr Tyr Lys			
	420	425	430			
	Ala Ile Thr Glu Ser	Val Arg Lys His Phe	Lys Gly Asn Gly Val Ile			
	435	440	445			
30	Ala Ser Met Glu Gln	Cys Asn Asp Phe Met	Leu Leu Gly Thr Glu Thr			
	450	455	460			
	Ile Cys Leu Gly Arg	Val Gly Asp Asp Phe	Trp Pro Thr Asp Pro Ser			
	465	470	475	480		
	Gly Asp Ile Asn Gly	Thr Tyr Trp Leu Gln	Gly Cys His Met Val His			
35	485	490	495			
	Cys Ala Tyr Asn Ser	Leu Trp Met Gly Asn	Phe Ile His Pro Asp Trp			
	500	505	510			
	Asp Met Phe Gln Ser	Thr His Pro Cys Ala	Glu Phe His Ala Ala Ser			
	515	520	525			
40	Arg Ala Ile Ser Gly	Gly Pro Ile Tyr Val	Ser Asp Val Val Gly Lys			

	530		535		540	
	His Asn Ile Pro Leu Leu Lys Arg Leu Val Leu Ala Asp Gly Ser Ile					
	545		550		555	560
	Leu Arg Cys Glu Tyr His Ala Leu Pro Thr Lys Asp Cys Leu Phe Val					
5		565		570		575
	Asp Pro Leu His Asp Gly Lys Thr Met Leu Lys Ile Trp Asn Leu Asn					
		580		585		590
	Lys Tyr Asn Gly Val Leu Gly Val Phe Asn Cys Gln Gly Gly Gly Trp					
		595		600		605
10	Ser Arg Glu Ser Arg Lys Asn Leu Cys Phe Ser Glu Tyr Ser Lys Pro					
	610		615		620	
	Ile Ser Cys Lys Thr Ser Pro Lys Asp Val Glu Trp Glu Asn Gly His					
	625		630		635	640
	Lys Pro Phe Pro Ile Lys Gly Val Glu Cys Phe Ala Met Tyr Phe Thr					
15		645		650		655
	Lys Glu Lys Lys Leu Ile Leu Ser Gln Leu Ser Asp Thr Ile Glu Ile					
		660		665		670
	Ser Leu Asp Pro Phe Asp Tyr Glu Leu Ile Val Val Ser Pro Met Thr					
		675		680		685
20	Ile Leu Pro Trp Glu Ser Ile Ala Phe Ala Pro Ile Gly Leu Val Asn					
	690		695		700	
	Met Leu Asn Ala Gly Gly Ala Val Lys Ser Leu Asp Ile Ser Glu Asp					
	705		710		715	720
	Asn Glu Asp Lys Met Val Gln Val Gly Ile Lys Gly Ala Gly Glu Met					
25		725		730		735
	Met Val Tyr Ser Ser Glu Lys Pro Lys Ala Cys Arg Val Asn Gly Glu					
		740		745		750
	Asp Met Glu Phe Glu Tyr Glu Glu Ser Met Ile Lys Val Gln Val Thr					
		755		760		765
30	Trp Asn His Asn Ser Gly Gly Phe Thr Thr Val Glu Tyr Leu Phe					
	770		775		780	783

	<210> 4
	<211> 2690
35	<212> DNA
	<213> Beta vulgaris L.
	<220>
	<221> CDS
40	<222> (236)... (2587)

<400> 4		
	ctaccaaat ccacaactta aagttcacct caatctttat tccatttttc ctcctaaac	60
	ttcattgtta agattttgta attgaattca aattcttaat tctgaatttt gtcatttttt	120
5	ttgtggggat atttataact atcatattat ttgtgtagat cattctacaa aaaagagagt	180
	gagttttttt agctcttatt tcctaagaaa ttaatagcaa aagttttgca taact atg	238
	Met	
	gct cca agc ttt agc aag gaa aat tcc aag acg tgt gat gag gtt gca	286
	Ala Pro Ser Phe Ser Lys Glu Asn Ser Lys Thr Cys Asp Glu Val Ala	
10	5 10 15	
	aac cat gat gat tgc aac acg tgt cca ata att tcc ttg gaa gaa tca	334
	Asn His Asp Asp Cys Asn Thr Cys Pro Ile Ile Ser Leu Glu Glu Ser	
	20 25 30	
	aac ttc atg gtg aat ggt cac gtg ata ttg tcc caa gtt cca tcc aac	382
15	Asn Phe Met Val Asn Gly His Val Ile Leu Ser Gln Val Pro Ser Asn	
	35 40 45	
	atc acg gcc att agt aaa atg ggt ttt gat ggg ctt ttt gtg ggt ttt	430
	Ile Thr Ala Ile Ser Lys Met Gly Phe Asp Gly Leu Phe Val Gly Phe	
	50 55 60 65	
20	gat gct cca gag ccc aag gcc cgg cac gtt gta tcc gtg ggc cag ctc	478
	Asp Ala Pro Glu Pro Lys Ala Arg His Val Val Ser Val Gly Gln Leu	
	70 75 80	
	aag gga att ccc ttc atg agt atc ttc agg ttc aag gta tgg tgg act	526
	Lys Gly Ile Pro Phe Met Ser Ile Phe Arg Phe Lys Val Trp Trp Thr	
25	85 90 95	
	acc cat tgg act ggg tcc aat ggg cgg gac ctt gag cat gag acc caa	574
	Thr His Trp Thr Gly Ser Asn Gly Arg Asp Leu Glu His Glu Thr Gln	
	100 105 110	
	att ctc atc ctt gat aag tca gat gaa ggt ttg ggc cgt ccc tat att	622
30	Ile Leu Ile Leu Asp Lys Ser Asp Glu Gly Leu Gly Arg Pro Tyr Ile	
	115 120 125	
	gtg atc ctc cca ttg atc gaa ggc cca ttt cgg gca tct ctc cag ccg	670
	Val Ile Leu Pro Leu Ile Glu Gly Pro Phe Arg Ala Ser Leu Gln Pro	
	130 135 140 145	
35	ggg tct gtt gat gac tat gtg gat ata tgt gtt gag agt ggg tcc act	718
	Gly Ser Val Asp Asp Tyr Val Asp Ile Cys Val Glu Ser Gly Ser Thr	
	150 155 160	
	aaa gtt gtc gga gac tcg ttc cgg gct gtt ctt tat ata cgg gct ggg	766
	Lys Val Val Gly Asp Ser Phe Arg Ala Val Leu Tyr Ile Arg Ala Gly	
40	165 170 175	

	cct gac cca ttt aag tta att aaa gat aca atg aag gaa gtc caa gcc	814
	Pro Asp Pro Phe Lys Leu Ile Lys Asp Thr Met Lys Glu Val Gln Ala	
	180 185 190	
5	cat tta ggg act ttc aaa ctc tta gat gac aaa act cct cca gga ata	862
	His Leu Gly Thr Phe Lys Leu Leu Asp Asp Lys Thr Pro Pro Gly Ile	
	195 200 205	
	gtg gac aag ttt gga tgg tgt aca tgg gat gca ttt tac ctc aaa gta	910
	Val Asp Lys Phe Gly Trp Cys Thr Trp Asp Ala Phe Tyr Leu Lys Val	
	210 215 220 225	
10	gag ccw tat ggt gtt tgg gaa gga gtt aaa gga ctc gtc gaa aac ggg	958
	Glu Pro Tyr Gly Val Trp Glu Gly Val Lys Gly Leu Val Glu Asn Gly	
	230 235 240	
	gtc cca ccc ggt ctc gta ctc att gat gat ggg tgg caa tct att tgt	1006
	Val Pro Pro Gly Leu Val Leu Ile Asp Asp Gly Trp Gln Ser Ile Cys	
15	245 250 255	
	cat gac gat gat ccg att acc gac caa gaa ggg ata aac cgg act tct	1054
	His Asp Asp Asp Pro Ile Thr Asp Gln Glu Gly Ile Asn Arg Thr Ser	
	260 265 270	
	gcc ggc gag caa atg cca tgt aga ttg atc aag tac gag gaa aac ttc	1102
20	Ala Gly Glu Gln Met Pro Cys Arg Leu Ile Lys Tyr Glu Glu Asn Phe	
	275 280 285	
	aag ttt agg gac tat aaa agc cca aat att atg ggc cat gag gat cat	1150
	Lys Phe Arg Asp Tyr Lys Ser Pro Asn Ile Met Gly His Glu Asp His	
	290 295 300 305	
25	ccc aat atg gga atg agg gcc ttt gtt agg gac ctt aag gag gag ttc	1198
	Pro Asn Met Gly Met Arg Ala Phe Val Arg Asp Leu Lys Glu Glu Phe	
	310 315 320	
	aaa act gtt gag cat gtg tat gtt tgg cat gct ttt acg ggc tat tgg	1246
	Lys Thr Val Glu His Val Tyr Val Trp His Ala Phe Thr Gly Tyr Trp	
30	325 330 335	
	gga ggg gta agg ccc aat gtt cca ggc cta ccr gag gcc caa gta gta	1294
	Gly Gly Val Arg Pro Asn Val Pro Gly Leu Pro Glu Ala Gln Val Val	
	340 345 350	
	acc cca aag ctt tcc ccg ggt ctt gag atg aca atg gaa gat cta gct	1342
35	Thr Pro Lys Leu Ser Pro Gly Leu Glu Met Thr Met Glu Asp Leu Ala	
	355 360 365	
	gtg gat aaa att gtt aat aat ggt att ggg ctt gtc cag cct gat aag	1390
	Val Asp Lys Ile Val Asn Asn Gly Ile Gly Leu Val Gln Pro Asp Lys	
	370 375 380 385	
40	gcc caa gaa ctt tat gaa ggg ttg cat tct cat ttg gaa aat tgt ggg	1438

	Ala Gln Glu Leu Tyr Glu Gly Leu His Ser His Leu Glu Asn Cys Gly	
	390 395 400	
	att gat gga gtc aaa gtt gat gtc atc cat ttg ttg gag atg atg gca	1486
	Ile Asp Gly Val Lys Val Asp Val Ile His Leu Leu Glu Met Met Ala	
5	405 410 415	
	gag gac tat gga gga aga gtt gaa cta gca aaa aca tac tat aag gca	1534
	Glu Asp Tyr Gly Gly Arg Val Glu Leu Ala Lys Thr Tyr Tyr Lys Ala	
	420 425 430	
	ata aca gaa tca gtg cgt aag cat ttc aaa ggc aac ggt gtg att gct	1582
10	Ile Thr Glu Ser Val Arg Lys His Phe Lys Gly Asn Gly Val Ile Ala	
	435 440 445	
	agc atg gag cag tgc aac gat ttc atg ctc ctt ggt act gag acc att	1630
	Ser Met Glu Gln Cys Asn Asp Phe Met Leu Leu Gly Thr Glu Thr Ile	
	450 455 460 465	
15	tgt ctt ggt cgc gtt ggg gat gac ttt tgg cca act gat ccg tct gga	1678
	Cys Leu Gly Arg Val Gly Asp Asp Phe Trp Pro Thr Asp Pro Ser Gly	
	470 475 480	
	gat ata aat ggt aca tat tgg ctc caa ggc tgt cat atg gtg cat tgt	1726
	Asp Ile Asn Gly Thr Tyr Trp Leu Gln Gly Cys His Met Val His Cys	
20	485 490 495	
	gcc tac aat agc tta tgg atg gga aac ttt ata cac cct gac tgg gac	1774
	Ala Tyr Asn Ser Leu Trp Met Gly Asn Phe Ile His Pro Asp Trp Asp	
	500 505 510	
	atg ttc caa tct aca cac cct tgt gct gaa ttt cat gct gca tct cgt	1822
25	Met Phe Gln Ser Thr His Pro Cys Ala Glu Phe His Ala Ala Ser Arg	
	515 520 525	
	gcg att tct ggt gga cca att tat gtt agt gat gtt gtt ggc aag cat	1870
	Ala Ile Ser Gly Gly Pro Ile Tyr Val Ser Asp Val Val Gly Lys His	
	530 535 540 545	
30	aac atc ccc ttg ctc aaa agg ctc gtc ttg gct gat ggt tcg atc ctt	1918
	Asn Ile Pro Leu Leu Lys Arg Leu Val Leu Ala Asp Gly Ser Ile Leu	
	550 555 560	
	cgt tgc gag tac cat gca ctt cct act aag gat tgc cta ttt gta gat	1966
	Arg Cys Glu Tyr His Ala Leu Pro Thr Lys Asp Cys Leu Phe Val Asp	
35	565 570 575	
	cct ttg cac gat ggc aaa aca atg ctc aaa att tgg aac ctc aac aag	2014
	Pro Leu His Asp Gly Lys Thr Met Leu Lys Ile Trp Asn Leu Asn Lys	
	580 585 590	
	tac aat gga gtg ctt gga gtc ttc aat tgc caa gga gga ggg tgg agc	2062
40	Tyr Asn Gly Val Leu Gly Val Phe Asn Cys Gln Gly Gly Gly Trp Ser	

	595	600	605	
	cgt gag tct cga aaa aat cta tgt ttc tca gag tat tca aaa cct att			2110
	Arg Glu Ser Arg Lys Asn Leu Cys Phe Ser Glu Tyr Ser Lys Pro Ile			
	610	615	620	625
5	tcc tgc aag aca agt cca aaa gat gtt gaa tgg gag aac gga cac aag			2158
	Ser Cys Lys Thr Ser Pro Lys Asp Val Glu Trp Glu Asn Gly His Lys			
	630	635	640	
	cca ttc ccc atc aaa gga gtg gaa tgt ttt gcc atg tac ttc acc aag			2206
	Pro Phe Pro Ile Lys Gly Val Glu Cys Phe Ala Met Tyr Phe Thr Lys			
10	645	650	655	
	gaa aaa aag cta atc ctc tca caa cta tct gac acc att gaa ata tca			2254
	Glu Lys Lys Leu Ile Leu Ser Gln Leu Ser Asp Thr Ile Glu Ile Ser			
	660	665	670	
15	ctt gat ccc ttc gat tac gag ctt att gta gtc tct ccg atg aca att			2302
	Leu Asp Pro Phe Asp Tyr Glu Leu Ile Val Val Ser Pro Met Thr Ile			
	675	680	685	
	cta ccc tgg gag tcg atc gca ttt gca ccc ata gga tta gta aac atg			2350
	Leu Pro Trp Glu Ser Ile Ala Phe Ala Pro Ile Gly Leu Val Asn Met			
	690	695	700	705
20	ctc aac gcc gga ggg gca gtc aag tct ttg gac atc agt gag gat aat			2398
	Leu Asn Ala Gly Gly Ala Val Lys Ser Leu Asp Ile Ser Glu Asp Asn			
	710	715	720	
	gag gat aag atg gtt cag gtt ggt att aaa ggg gcc gga gaa atg atg			2446
	Glu Asp Lys Met Val Gln Val Gly Ile Lys Gly Ala Gly Glu Met Met			
25	725	730	735	
	gtt tat tca tca gaa aag cca aaa gcg tgt aga gtt aat gga gaa gac			2494
	Val Tyr Ser Ser Glu Lys Pro Lys Ala Cys Arg Val Asn Gly Glu Asp			
	740	745	750	
	atg gag ttt gag tat gaa gag agc atg att aag gtt caa gtt aca tgg			2542
30	Met Glu Phe Glu Tyr Glu Glu Ser Met Ile Lys Val Gln Val Thr Trp			
	755	760	765	
	aac cat aac tca ggt ggt ttt acc act gtt gag tac tta ttt tga gcttg			2592
	Asn His Asn Ser Gly Gly Phe Thr Thr Val Glu Tyr Leu Phe			
	770	775	780	
35	aagctaactct aagtctttac ttaatgagtg atgtaactga gtagttgact tgagagtaca			2652
	gtatgtgtga agcttattat tccaaaaaaa aaaaaaaa			2690
	<210> 5			
	<211> 777			
40	<212> PRT			

<213> Brassica juncea

<400> 5

	Met	Ala	Pro	Pro	Ser	Val	Ile	Lys	Ser	Asp	Ala	Ala	Val	Asn	Gly	Ile
						5				10					15	
5	Asp	Leu	Ser	Gly	Lys	Pro	Leu	Phe	Arg	Leu	Glu	Gly	Ser	Asp	Leu	Leu
				20					25					30		
	Ala	Asn	Gly	His	Val	Val	Leu	Thr	Asp	Val	Pro	Val	Asn	Val	Thr	Val
			35					40					45			
10	Thr	Ala	Ser	Pro	Tyr	Leu	Ala	Asp	Lys	Asp	Gly	Glu	Pro	Val	Asp	Ala
		50					55				60					
	Ser	Ala	Gly	Ser	Phe	Ile	Gly	Phe	Asn	Leu	Asp	Gly	Glu	Pro	Arg	Ser
	65					70					75				80	
	Arg	His	Val	Ala	Ser	Ile	Gly	Lys	Leu	Arg	Asp	Ile	Arg	Phe	Met	Ser
15					85					90					95	
	Ile	Phe	Arg	Phe	Lys	Val	Trp	Trp	Thr	Thr	His	Trp	Val	Gly	Ser	Lys
				100					105					110		
	Gly	Ser	Asp	Ile	Glu	Asn	Glu	Thr	Gln	Ile	Ile	Ile	Leu	Glu	Asn	Ser
			115					120					125			
20	Gly	Ser	Gly	Arg	Pro	Tyr	Val	Leu	Leu	Leu	Pro	Leu	Leu	Glu	Gly	Ser
		130					135				140					
	Phe	Arg	Ser	Ser	Phe	Gln	Pro	Gly	Glu	Asp	Asp	Asp	Val	Ala	Val	Cys
	145					150					155				160	
	Val	Glu	Ser	Gly	Ser	Thr	Gln	Val	Thr	Gly	Ser	Glu	Phe	Arg	Gln	Val
25				165						170				175		
	Val	Tyr	Val	His	Ala	Gly	Asp	Asp	Pro	Phe	Lys	Leu	Val	Lys	Asp	Ala
				180					185					190		
	Met	Lys	Val	Val	Arg	Val	His	Met	Asn	Thr	Phe	Lys	Leu	Leu	Glu	Glu
		195						200					205			
30	Lys	Thr	Pro	Pro	Gly	Ile	Val	Asp	Lys	Phe	Gly	Trp	Cys	Thr	Trp	Asp
		210					215					220				
	Ala	Phe	Tyr	Leu	Thr	Val	Asn	Pro	Asp	Gly	Val	His	Lys	Gly	Val	Lys
	225					230				235					240	
	Cys	Leu	Val	Asp	Gly	Gly	Cys	Pro	Pro	Gly	Leu	Val	Leu	Ile	Asp	Asp
35				245					250					255		
	Gly	Trp	Gln	Ser	Ile	Gly	His	Asp	Ser	Asp	Gly	Ile	Asp	Val	Glu	Gly
			260					265					270			
	Met	Ser	Cys	Thr	Val	Ala	Gly	Glu	Gln	Met	Pro	Cys	Arg	Leu	Leu	Lys
		275						280					285			
40	Phe	Gln	Glu	Asn	Phe	Lys	Phe	Arg	Asp	Tyr	Val	Ser	Pro	Lys	Asp	Lys

	290		295		300
	Asn Glu Val Gly Met Lys Ala Phe Val Arg Asp Leu Lys Glu Glu Phe				
	305		310		315
	Ser Thr Val Asp Tyr Ile Tyr Val Trp His Ala Leu Cys Gly Tyr Trp				
5		325		330	335
	Gly Gly Leu Arg Pro Gly Ala Pro Thr Leu Pro Pro Ser Thr Ile Val				
		340		345	350
	Arg Pro Glu Leu Ser Pro Gly Leu Lys Leu Thr Met Gln Asp Leu Ala				
		355		360	365
10	Val Asp Lys Ile Val Asp Thr Gly Ile Gly Phe Val Ser Pro Asp Met				
		370		375	380
	Ala Asn Glu Phe Tyr Glu Gly Leu His Ser His Leu Gln Asn Val Gly				
	385		390		395
	Ile Asp Gly Val Lys Val Asp Val Ile His Ile Leu Glu Met Leu Cys				
15		405		410	415
	Glu Lys Tyr Gly Gly Arg Val Asp Leu Ala Lys Ala Tyr Phe Lys Ala				
		420		425	430
	Leu Thr Ser Ser Val Asn Lys His Phe Asp Gly Asn Gly Val Ile Ala				
		435		440	445
20	Ser Met Glu His Cys Asn Asp Phe Met Phe Leu Gly Thr Glu Ala Ile				
		450		455	460
	Ser Leu Gly Arg Val Gly Asp Asp Phe Trp Cys Thr Asp Pro Ser Gly				
	465		470		475
	Asp Ile Asn Gly Thr Tyr Trp Leu Gln Gly Cys His Met Val His Cys				
25		485		490	495
	Ala Tyr Asn Ser Leu Trp Met Gly Asn Phe Ile Gln Pro Asp Trp Asp				
		500		505	510
	Met Phe Gln Ser Thr His Pro Cys Ala Glu Phe His Ala Ala Ser Arg				
		515		520	525
30	Ala Ile Ser Gly Gly Pro Ile Tyr Ile Ser Asp Cys Val Gly Gln His				
		530		535	540
	Asp Phe Asp Leu Leu Lys Arg Leu Val Leu Pro Asp Gly Ser Ile Leu				
	545		550		555
	Arg Cys Glu His Tyr Ala Leu Pro Thr Arg Asp Arg Leu Phe Glu Asp				
35		565		570	575
	Pro Leu His Asp Gly Lys Thr Met Leu Lys Ile Trp Asn Leu Asn Lys				
		580		585	590
	Tyr Thr Gly Ile Ile Gly Ala Phe Asn Cys Gln Gly Gly Gly Trp Cys				
		595		600	605
40	Arg Glu Thr Arg Arg Asn Gln Cys Phe Ser Gln Cys Val Asn Thr Leu				

	610	615	620	
	Thr Ala Thr Thr Asn Pro Lys Asp Val Glu Trp Asn Ser Gly Asn Asn			
	625	630	635	640
	Pro Ile Ser Val Glu Asn Val Glu Glu Phe Ala Leu Phe Leu Ser Gln			
5	645	650	655	
	Ser Lys Lys Leu Val Leu Ser Gly Pro Asn Asp Asp Leu Glu Ile Thr			
	660	665	670	
	Leu Glu Pro Phe Lys Phe Glu Leu Ile Thr Val Ser Pro Val Val Thr			
	675	680	685	
10	Ile Glu Gly Ser Ser Val Gln Phe Ala Pro Ile Gly Leu Val Asn Met			
	690	695	700	
	Leu Asn Thr Ser Gly Ala Ile Arg Ser Leu Val Tyr His Glu Glu Ser			
	705	710	715	720
	Val Glu Ile Gly Val Arg Gly Ala Gly Glu Phe Arg Val Tyr Ala Ser			
15	725	730	735	
	Arg Lys Pro Ala Ser Cys Lys Ile Asp Gly Glu Val Val Glu Phe Gly			
	740	745	750	
	Tyr Glu Glu Ser Met Val Met Val Gln Val Pro Trp Ser Ala Pro Glu			
	755	760	765	
20	Gly Leu Ser Ser Ile Lys Tyr Glu Phe			
	770	775	777	
	<210> 6			
	<211> 2690			
25	<212> DNA			
	<213> Brassica juncea			
	<220>			
	<221> CDS			
30	<222> (134)... (2467)			
	<400> 6			
	accaatccaa aatctcatca aataatcgca attaggggaa gtttacaaga ttcacatct			60
	cggttactat ataactacgc tcttcttcc tgcctaate caacttaacc taaaaaccac			120
35	tctatcagcg aaa atg gct cca ccg agc gta att aaa tcc gat gct gca			169
	Met Ala Pro Pro Ser Val Ile Lys Ser Asp Ala Ala			
		5	10	
	gtc aac ggc att gac ctc tcc gga aag ccg ctt ttc cgg cta gag ggt			217
	Val Asn Gly Ile Asp Leu Ser Gly Lys Pro Leu Phe Arg Leu Glu Gly			
40	15	20	25	

	tcc gat ctc cta gcc aat ggt cac gtt gtc tta acc gat gta ccg gtt	265
	Ser Asp Leu Leu Ala Asn Gly His Val Val Leu Thr Asp Val Pro Val	
	30 35 40	
5	aac gtg act gtc act gct tca cct tac cta gct gac aaa gac gga gaa	313
	Asn Val Thr Val Thr Ala Ser Pro Tyr Leu Ala Asp Lys Asp Gly Glu	
	45 50 55 60	
	ccg gtt gac gcc tcc gct ggt tca ttc atc ggg ttt aat ctc gac ggt	361
	Pro Val Asp Ala Ser Ala Gly Ser Phe Ile Gly Phe Asn Leu Asp Gly	
	65 70 75	
10	gag cca cga agc cgc cac gtg gcg tcc atc ggt aaa ctc agg gat att	409
	Glu Pro Arg Ser Arg His Val Ala Ser Ile Gly Lys Leu Arg Asp Ile	
	80 85 90	
	cga ttc atg agc ata ttc cgt ttc aag gtt tgg tgg act act cac tgg	457
	Arg Phe Met Ser Ile Phe Arg Phe Lys Val Trp Trp Thr Thr His Trp	
15	95 100 105	
	gtc ggt tcc aaa gga tcc gac atc gag aac gag acc cag atc atc atc	505
	Val Gly Ser Lys Gly Ser Asp Ile Glu Asn Glu Thr Gln Ile Ile Ile	
	110 115 120	
20	ctc gag aac tcc ggg tcg ggt cgt cct tat gtt ctt ctt ctg ccg ctt	553
	Leu Glu Asn Ser Gly Ser Gly Arg Pro Tyr Val Leu Leu Leu Pro Leu	
	125 130 135 140	
	ctt gaa ggc tct ttc cgt tca tcc ttt cag cct ggg gaa gac gat gac	601
	Leu Glu Gly Ser Phe Arg Ser Ser Phe Gln Pro Gly Glu Asp Asp Asp	
	145 150 155	
25	gtg gcg gtt tgt gtc gaa tcc ggg tcg acc cag gtg acc ggg tcg gag	649
	Val Ala Val Cys Val Glu Ser Gly Ser Thr Gln Val Thr Gly Ser Glu	
	160 165 170	
	ttt cgt caa gtt gtg tat gtt cac gcc gga gac gat ccg ttc aag ctc	697
	Phe Arg Gln Val Val Tyr Val His Ala Gly Asp Asp Pro Phe Lys Leu	
30	175 180 185	
	gtg aaa gac gcg atg aag gtg gtt agg gtt cat atg aac acc ttc aag	745
	Val Lys Asp Ala Met Lys Val Val Arg Val His Met Asn Thr Phe Lys	
	190 195 200	
35	ctc ttg gaa gag aag acr ccg ccg gga atc gtc gat aag ttc ggg tgg	793
	Leu Leu Glu Glu Lys Thr Pro Pro Gly Ile Val Asp Lys Phe Gly Trp	
	205 210 215 220	
	tgc acg tgg gat gcg ttt tat ttg acg gtg aac cct gac gga gtt cat	841
	Cys Thr Trp Asp Ala Phe Tyr Leu Thr Val Asn Pro Asp Gly Val His	
	225 230 235	
40	aag ggt gtt aag tgt ctc gtc gac ggt ggt tgt ccg ccg gga ttg gtc	889

	Lys Gly Val Lys Cys Leu Val Asp Gly Gly Cys Pro Pro Gly Leu Val	
	240 245 250	
	cta atc gac gac ggt tgg caa tcg att gga cat gac tcc gat ggt atc	937
	Leu Ile Asp Asp Gly Trp Gln Ser Ile Gly His Asp Ser Asp Gly Ile	
5	255 260 265	
	gat gtt gaa ggg atg agt tgt acc gtc gcc ggg gag caa atg cct tgc	985
	Asp Val Glu Gly Met Ser Cys Thr Val Ala Gly Glu Gln Met Pro Cys	
	270 275 280	
	agg ctt ctg aaa ttt caa gag aac ttc aag ttc aga gac tac gtc tct	1033
10	Arg Leu Leu Lys Phe Gln Glu Asn Phe Lys Phe Arg Asp Tyr Val Ser	
	285 290 295 300	
	ccg aaa gac aaa aac gaa gtc ggg atg aaa gct ttc gtc aga gat ctg	1081
	Pro Lys Asp Lys Asn Glu Val Gly Met Lys Ala Phe Val Arg Asp Leu	
	305 310 315	
15	aaa gaa gaa ttc tcc acc gtt gat tac atc tac gtc tgg cac gcg ctt	1129
	Lys Glu Glu Phe Ser Thr Val Asp Tyr Ile Tyr Val Trp His Ala Leu	
	320 325 330	
	tgc ggc tac tgg ggt ggt ctt cgt ccc gga gct cct act ctt ccg ccc	1177
	Cys Gly Tyr Trp Gly Gly Leu Arg Pro Gly Ala Pro Thr Leu Pro Pro	
20	335 340 345	
	tca act att gtc cgg cca gag ctc tcg ccg ggg ctt aag ttg acg atg	1225
	Ser Thr Ile Val Arg Pro Glu Leu Ser Pro Gly Leu Lys Leu Thr Met	
	350 355 360	
	caa gat ctc gcc gtt gat aag att gtc gat acc gga atc gga ttc gtc	1273
25	Gln Asp Leu Ala Val Asp Lys Ile Val Asp Thr Gly Ile Gly Phe Val	
	365 370 375 380	
	tcg ccg gac atg gcg aat gag ttt tac gaa ggt ctt cac tct cat ctt	1321
	Ser Pro Asp Met Ala Asn Glu Phe Tyr Glu Gly Leu His Ser His Leu	
	385 390 395	
30	caa aac gtc ggt att gac ggc gtt aaa gtt gac gtc atc cac ata ttg	1369
	Gln Asn Val Gly Ile Asp Gly Val Lys Val Asp Val Ile His Ile Leu	
	400 405 410	
	gag atg ttg tgc gag aaa tat ggc ggg aga gta gac ttg gct aaa gct	1417
	Glu Met Leu Cys Glu Lys Tyr Gly Gly Arg Val Asp Leu Ala Lys Ala	
35	415 420 425	
	tac ttc aag gcg tta act tcc tca gtg aat aag cat ttt gac ggt aac	1465
	Tyr Phe Lys Ala Leu Thr Ser Ser Val Asn Lys His Phe Asp Gly Asn	
	430 435 440	
	ggc gtt atc gct agc atg gag cac tgt aat gat ttc atg ttc ctt gga	1513
40	Gly Val Ile Ala Ser Met Glu His Cys Asn Asp Phe Met Phe Leu Gly	

	445	450	455	460	
	acc gaa gcc atc tct cta ggt cgt gtc ggt gat gac ttt tgg tgc acg				1561
	Thr Glu Ala Ile Ser Leu Gly Arg Val Gly Asp Asp Phe Trp Cys Thr				
		465	470	475	
5	gat cca tca ggc gac ata aac ggc aca tat tgg ctg caa gga tgc cac				1609
	Asp Pro Ser Gly Asp Ile Asn Gly Thr Tyr Trp Leu Gln Gly Cys His				
		480	485	490	
	atg gtc cac tgt gcc tac aac agt ctt tgg atg gga aat ttc atc cag				1657
	Met Val His Cys Ala Tyr Asn Ser Leu Trp Met Gly Asn Phe Ile Gln				
10		495	500	505	
	cct gat tgg gac atg ttt cag tcc aca cat cct tgt gct gag ttc cat				1705
	Pro Asp Trp Asp Met Phe Gln Ser Thr His Pro Cys Ala Glu Phe His				
		510	515	520	
	gct gct tct cgt gcc atc tcc ggt ggg ccc att tac atc agc gat tgt				1753
15	Ala Ala Ser Arg Ala Ile Ser Gly Gly Pro Ile Tyr Ile Ser Asp Cys				
		525	530	535	540
	gtg ggc cag cac gat ttc gat ctc ttg aag cga ctc gtc ttg cct gac				1801
	Val Gly Gln His Asp Phe Asp Leu Leu Lys Arg Leu Val Leu Pro Asp				
		545	550	555	
20	ggt tcg att ttg agg tgt gag cac tat gca ctc cca act cgt gac cgt				1849
	Gly Ser Ile Leu Arg Cys Glu His Tyr Ala Leu Pro Thr Arg Asp Arg				
		560	565	570	
	ctc ttt gaa gac cct ctt cat gat ggc aaa acc atg ctc aag att tgg				1897
	Leu Phe Glu Asp Pro Leu His Asp Gly Lys Thr Met Leu Lys Ile Trp				
25		575	580	585	
	aac ttg aac aag tac act gga att att gga gca ttc aac tgc caa gga				1945
	Asn Leu Asn Lys Tyr Thr Gly Ile Ile Gly Ala Phe Asn Cys Gln Gly				
		590	595	600	
	gga gga tgg tgc aga gaa acc cga cgc aac caa tgc ttc tcc caa tgc				1993
30	Gly Gly Trp Cys Arg Glu Thr Arg Arg Asn Gln Cys Phe Ser Gln Cys				
		605	610	615	620
	gtt aac acg tta acc gcc aca aca aat cct aag gac gtt gaa tgg aac				2041
	Val Asn Thr Leu Thr Ala Thr Thr Asn Pro Lys Asp Val Glu Trp Asn				
		625	630	635	
35	agt ggg aac aac cca atc tcc gtt gaa aac gtt gaa gag ttt gct ttg				2089
	Ser Gly Asn Asn Pro Ile Ser Val Glu Asn Val Glu Glu Phe Ala Leu				
		640	645	650	
	ttc ttg tct cag tct aag aag ctt gtg ttg tct gga cca aac gat gat				2137
	Phe Leu Ser Gln Ser Lys Lys Leu Val Leu Ser Gly Pro Asn Asp Asp				
40		655	660	665	

	ctc gag atc act ttg gag cct ttc aag ttt gag cta atc act gtc tca	2185
	Leu Glu Ile Thr Leu Glu Pro Phe Lys Phe Glu Leu Ile Thr Val Ser	
	670 675 680	
5	cca gtt gtc act att gag ggt agt tcg gtt cag ttt gct cca atc gga	2233
	Pro Val Val Thr Ile Glu Gly Ser Ser Val Gln Phe Ala Pro Ile Gly	
	685 690 695 700	
	ttg gtt aac atg cta aac act agc ggt gca att cga tcc ttg gtg tat	2281
	Leu Val Asn Met Leu Asn Thr Ser Gly Ala Ile Arg Ser Leu Val Tyr	
	705 710 715	
10	cat gag gaa tcc gtt gag att gga gtt cgt ggt gct gga gag ttc agg	2329
	His Glu Glu Ser Val Glu Ile Gly Val Arg Gly Ala Gly Glu Phe Arg	
	720 725 730	
	gtt tat gca tca agg aaa cct gcg agc tgc aaa att gat ggt gaa gtt	2377
	Val Tyr Ala Ser Arg Lys Pro Ala Ser Cys Lys Ile Asp Gly Glu Val	
15	735 740 745	
	gtt gag ttt gga tac gaa gag tca atg gtg atg gtt caa gtg cct tgg	2425
	Val Glu Phe Gly Tyr Glu Glu Ser Met Val Met Val Gln Val Pro Trp	
	750 755 760	
20	tct gca ccc gag ggt ttg tct tct att aag tat gag ttt tag agtttccga	2476
	Ser Ala Pro Glu Gly Leu Ser Ser Ile Lys Tyr Glu Phe	
	765 770 775	
	agggtcttat ttgtatcctt ctaaactcct taattatgag ctccgtgccg tttctttttc	2536
	tatatggttt ctgagagtga acatctaata tttaccact agggataat tattggcttt	2596
	taagtgattt gtttttgaac tgtttttagt ggtgtaattt gtactgcccc tattattttt	2656
25	catattttatt tgtgaaagat aaaaaaaaaa aaaa	2690
	<210> 7	
	<211> 572	
	<212> PRT	
30	<213> Brassica napus	
	<400> 7	
	Leu Glu Glu Lys Thr Pro Pro Gly Ile Val Asp Lys Phe Gly Trp Cys	
	5 10 15	
35	Thr Trp Asp Ala Phe Tyr Leu Thr Val Asn Pro Asp Gly Val His Lys	
	20 25 30	
	Gly Val Lys Cys Leu Val Asp Gly Cys Pro Pro Gly Leu Val Leu	
	35 40 45	
40	Ile Asp Asp Gly Trp Gln Ser Ile Gly His Asp Ser Asp Gly Ile Asp	
	50 55 60	

	Val	Glu	Gly	Met	Ser	Cys	Thr	Val	Ala	Gly	Glu	Gln	Met	Pro	Cys	Arg	
	65					70					75					80	
	Leu	Pro	Lys	Phe	Gln	Glu	Asn	Phe	Lys	Phe	Arg	Asp	Tyr	Val	Ser	Pro	
					85					90					95		
5	Lys	Asp	Lys	Asn	Glu	Val	Gly	Met	Lys	Ala	Phe	Val	Arg	Asp	Leu	Lys	
				100				105						110			
	Glu	Glu	Phe	Ser	Thr	Val	Asp	Tyr	Ile	Tyr	Val	Trp	His	Ala	Leu	Cys	
			115				120						125				
	Gly	Tyr	Trp	Gly	Gly	Leu	Arg	Pro	Gly	Ala	Pro	Thr	Leu	Pro	Pro	Ser	
10		130				135					140						
	Thr	Ile	Val	Arg	Pro	Glu	Leu	Ser	Pro	Gly	Leu	Lys	Leu	Thr	Met	Gln	
	145				150					155						160	
	Asp	Leu	Ala	Val	Asp	Lys	Ile	Ile	Asp	Thr	Gly	Ile	Gly	Phe	Val	Ser	
				165					170						175		
15	Pro	Asp	Met	Ala	Asn	Glu	Phe	Tyr	Glu	Gly	Leu	His	Ser	His	Leu	Gln	
			180					185					190				
	Asn	Val	Gly	Ile	Asn	Gly	Val	Lys	Val	Asp	Val	Ile	His	Ile	Leu	Glu	
		195				200						205					
	Met	Leu	Cys	Glu	Lys	Tyr	Gly	Gly	Arg	Val	Asp	Leu	Ala	Lys	Ala	Tyr	
20		210				215					220						
	Phe	Lys	Ala	Leu	Thr	Ser	Ser	Val	Asn	Lys	His	Phe	Asp	Gly	Asn	Ala	
	225				230					235						240	
	Val	Ile	Ala	Ser	Met	Glu	His	Cys	Asn	Asp	Phe	Met	Phe	Leu	Gly	Thr	
				245					250						255		
25	Glu	Ala	Ile	Ser	Leu	Gly	Arg	Val	Gly	Asp	Asp	Phe	Trp	Cys	Thr	Asp	
		260						265					270				
	Pro	Ser	Gly	Asp	Ile	Asn	Gly	Thr	Tyr	Trp	Leu	Gln	Gly	Cys	His	Met	
		275					280						285				
	Val	His	Cys	Ala	Tyr	Asn	Ser	Leu	Trp	Met	Gly	Asn	Phe	Ile	Gln	Pro	
30		290				295						300					
	Asp	Trp	Asp	Met	Phe	Gln	Ser	Thr	His	Pro	Cys	Ala	Glu	Phe	His	Ala	
	305				310					315						320	
	Ala	Ser	Arg	Ala	Ile	Ser	Gly	Gly	Pro	Ile	Tyr	Ile	Ser	Asp	Cys	Val	
				325					330						335		
35	Gly	Gln	His	Asp	Phe	Asp	Leu	Leu	Arg	Arg	Leu	Val	Leu	Pro	Asp	Gly	
		340						345					350				
	Ser	Ile	Leu	Arg	Cys	Glu	Tyr	Tyr	Ala	Leu	Pro	Thr	Arg	Asp	Arg	Leu	
		355					360						365				
	Phe	Glu	Asp	Pro	Leu	His	Asp	Gly	Lys	Thr	Met	Leu	Lys	Ile	Trp	Asn	
40		370				375						380					

	Leu Asn Lys Tyr Thr Gly Ile Ile Gly Ala Phe Asn Cys Gln Gly Gly	
	385 390 395 400	
	Gly Trp Cys Arg Glu Thr Arg Arg Asp Gln Cys Phe Ser Gln Cys Val	
	405 410 415	
5	Asn Thr Leu Thr Ala Thr Thr Asn Pro Asn Asp Val Glu Trp Asn Ser	
	420 425 430	
	Gly Asn Asn Pro Ile Ser Ile Glu Asn Val Glu Glu Phe Ala Leu Phe	
	435 440 445	
	Leu Ser Gln Ser Lys Lys Leu Val Leu Ser Gly Gln Asn Asp Asp Leu	
10	450 455 460	
	Glu Ile Thr Leu Glu Pro Phe Lys Phe Glu Leu Ile Thr Val Ser Pro	
	465 470 475 480	
	Val Val Thr Ile Glu Gly Ser Ser Val Gln Phe Ala Pro Ile Gly Leu	
	485 490 495	
15	Val Asn Met Leu Asn Thr Ser Gly Ala Ile Arg Ser Leu Val Tyr His	
	500 505 510	
	Glu Glu Ser Val Glu Ile Gly Val Arg Gly Ala Gly Glu Phe Arg Val	
	515 520 525	
	Tyr Ala Ser Lys Lys Pro Val Ser Cys Lys Ile Asp Gly Glu Asp Val	
20	530 535 540	
	Glu Phe Gly Tyr Glu Glu Ser Met Val Met Val Gln Val Pro Trp Ser	
	545 550 555 560	
	Ala Pro Glu Gly Leu Ser Ser Ile Lys Tyr Leu Phe	
	565 570 572	
25		
	<210> 8	
	<211> 1762	
	<212> DNA	
	<213> Brassica napus	
30		
	<220>	
	<221> CDS	
	<222> (1)... (1719)	
35		
	<400> 8	
	ttg gaa gaa aaa acg ccg ccg gga atc gtc gat aag ttc ggg tgg tgc	48
	Leu Glu Glu Lys Thr Pro Pro Gly Ile Val Asp Lys Phe Gly Trp Cys	
	5 10 15	
	acg tgg gat gcg ttt tat ttg acg gtg aac cct gac gga gtt cat aag	96
40	Thr Trp Asp Ala Phe Tyr Leu Thr Val Asn Pro Asp Gly Val His Lys	

		20		25		30		
		ggt gtt aag tgt ctc gtc gac ggt ggt tgt ccg ccg gga ttg gtc cta	144					
		Gly Val Lys Cys Leu Val Asp Gly Gly Cys Pro Pro Gly Leu Val Leu						
		35		40		45		
5		atc gac gac ggt tgg caa tcg att gga cat gac tcc gat ggt atc gat	192					
		Ile Asp Asp Gly Trp Gln Ser Ile Gly His Asp Ser Asp Gly Ile Asp						
		50		55		60		
		ggt gaa ggg atg agt tgt acc gtc gcc ggg gag caa atg cct tgc agg	240					
		Val Glu Gly Met Ser Cys Thr Val Ala Gly Glu Gln Met Pro Cys Arg						
10		65		70		75		80
		ctt ccg aaa ttt caa gag aac ttc aag ttc aga gac tac gtc tct ccg	288					
		Leu Pro Lys Phe Gln Glu Asn Phe Lys Phe Arg Asp Tyr Val Ser Pro						
		85		90		95		
		aaa gac aaa aac gaa gtc ggg atg aaa gct ttc gtc aga gat ctg aaa	336					
15		Lys Asp Lys Asn Glu Val Gly Met Lys Ala Phe Val Arg Asp Leu Lys						
		100		105		110		
		gaa gaa ttc tcc acc gtt gat tac atc tac gtc tgg cac gcg ctt tgc	384					
		Glu Glu Phe Ser Thr Val Asp Tyr Ile Tyr Val Trp His Ala Leu Cys						
		115		120		125		
20		ggy tac tgg gdw ggt ctt cgt ccc gga gct cct act ctt ccg ccs tcr	432					
		Gly Tyr Trp Gly Gly Leu Arg Pro Gly Ala Pro Thr Leu Pro Pro Ser						
		130		135		140		
		act att gtc cgr cca gag ctc tcg ccg ggg ctt aag ttg acg atg caa	480					
		Thr Ile Val Arg Pro Glu Leu Ser Pro Gly Leu Lys Leu Thr Met Gln						
25		145		150		155		160
		gat ctc gcc gtt gat aag atc atc gat acc gga atc gga ttc gtc tcg	528					
		Asp Leu Ala Val Asp Lys Ile Ile Asp Thr Gly Ile Gly Phe Val Ser						
		165		170		175		
		ccg gac atg gcg aac gag ttt tac gaa ggt ctt cac tct cat ctt caa	576					
30		Pro Asp Met Ala Asn Glu Phe Tyr Glu Gly Leu His Ser His Leu Gln						
		180		185		190		
		aac gtc ggc att aac ggc gtt aaa gtt gac gtt atc cac ata ctg gag	624					
		Asn Val Gly Ile Asn Gly Val Lys Val Asp Val Ile His Ile Leu Glu						
		195		200		205		
35		atg ttg tgc gag aaa tat ggc ggg aga gtt gac ttg gct aaa gct tac	672					
		Met Leu Cys Glu Lys Tyr Gly Gly Arg Val Asp Leu Ala Lys Ala Tyr						
		210		215		220		
		ttc aag gcg tta acg tcg tca gtg aat aag cat ttt gac ggc aac gcc	720					
		Phe Lys Ala Leu Thr Ser Ser Val Asn Lys His Phe Asp Gly Asn Ala						
40		225		230		235		240

	gtt atc gcc agc atg gag cac tgt aat gac ttc atg ttc ctt gga acc	768
	Val Ile Ala Ser Met Glu His Cys Asn Asp Phe Met Phe Leu Gly Thr	
	245 250 255	
5	gaa gcc atc tct cta ggt cgt gtc ggt gat gac ttt tgg tgc acg gat	816
	Glu Ala Ile Ser Leu Gly Arg Val Gly Asp Asp Phe Trp Cys Thr Asp	
	260 265 270	
	cca tct ggc gac att aac ggc acg tat tgg ctg caa gga tgt cac atg	864
	Pro Ser Gly Asp Ile Asn Gly Thr Tyr Trp Leu Gln Gly Cys His Met	
	275 280 285	
10	gtc cac tgt gcc tac aac agt ctt tgg atg gga aat ttc atc cag cct	912
	Val His Cys Ala Tyr Asn Ser Leu Trp Met Gly Asn Phe Ile Gln Pro	
	290 295 300	
	gat tgg gac atg ttt cag tcc aca cat cct tgt gct gag ttc cat gct	960
	Asp Trp Asp Met Phe Gln Ser Thr His Pro Cys Ala Glu Phe His Ala	
15	305 310 315 320	
	gct tca cgt gcc atc tcc ggt ggg ccc att tac atc agc gat tgt gtg	1008
	Ala Ser Arg Ala Ile Ser Gly Gly Pro Ile Tyr Ile Ser Asp Cys Val	
	325 330 335	
	ggc cag cac gat ttc gat ctc ttg agg aga ctc gtt ttg cct gac ggt	1056
20	Gly Gln His Asp Phe Asp Leu Leu Arg Arg Leu Val Leu Pro Asp Gly	
	340 345 350	
	tcg att ttg agg tgt gag tac tat gct ctc cca act cgt gac cgt ctc	1104
	Ser Ile Leu Arg Cys Glu Tyr Tyr Ala Leu Pro Thr Arg Asp Arg Leu	
	355 360 365	
25	ttt gaa gac cct ctt cat gat ggc aaa acc atg ctc aag att tgg aac	1152
	Phe Glu Asp Pro Leu His Asp Gly Lys Thr Met Leu Lys Ile Trp Asn	
	370 375 380	
	ttg aac aag tac act gga atc atc gga gca ttc aac tgt caa gga gga	1200
	Leu Asn Lys Tyr Thr Gly Ile Ile Gly Ala Phe Asn Cys Gln Gly Gly	
30	385 390 395 400	
	gga tgg tgc aga gaa act cga cgc gac caa tgc ttc tcc caa tgc gtt	1248
	Gly Trp Cys Arg Glu Thr Arg Arg Asp Gln Cys Phe Ser Gln Cys Val	
	405 410 415	
	aac acg tta acc gcc aca aca aat cct aat gac gtt gaa tgg aac agt	1296
35	Asn Thr Leu Thr Ala Thr Thr Asn Pro Asn Asp Val Glu Trp Asn Ser	
	420 425 430	
	ggg aac aac ccg atc tcc att gaa aac gtt gaa gag ttt gct ttg ttc	1344
	Gly Asn Asn Pro Ile Ser Ile Glu Asn Val Glu Glu Phe Ala Leu Phe	
	435 440 445	
40	ttg tct caa tcc aag aag ctt gtg ttg tcc ggg caa aac gat gat ctc	1392

	Leu Ser Gln Ser Lys Lys Leu Val Leu Ser Gly Gln Asn Asp Asp Leu	
	450 455 460	
	gag atc aca tta gag ccc ttc aag ttc gag ctc atc act gtc tca cca	1440
	Glu Ile Thr Leu Glu Pro Phe Lys Phe Glu Leu Ile Thr Val Ser Pro	
5	465 470 475 480	
	gtt gtc acc att gag ggc agt tcg gtt cag ttt gct cca atc gga ttg	1488
	Val Val Thr Ile Glu Gly Ser Ser Val Gln Phe Ala Pro Ile Gly Leu	
	485 490 495	
	gtt aac atg ctt aac act agc ggt gcg att cga tcc ttg gtt tat cat	1536
10	Val Asn Met Leu Asn Thr Ser Gly Ala Ile Arg Ser Leu Val Tyr His	
	500 505 510	
	gag gaa tcc gtt gag atc ggt gtt cgt ggt gct gga gaa ttc agg gtt	1584
	Glu Glu Ser Val Glu Ile Gly Val Arg Gly Ala Gly Glu Phe Arg Val	
	515 520 525	
15	tat gca tcg aag aaa cct gtg agc tgc aag att gat ggt gaa gat gtt	1632
	Tyr Ala Ser Lys Lys Pro Val Ser Cys Lys Ile Asp Gly Glu Asp Val	
	530 535 540	
	gag ttt ggg tac gaa gag tca atg gtg atg gtt caa gtg cct tgg tct	1680
	Glu Phe Gly Tyr Glu Glu Ser Met Val Met Val Gln Val Pro Trp Ser	
20	545 550 555 560	
	gca cca gag ggt ttg tct tct att aag tat ttg ttt tag agttatttaa	1729
	Ala Pro Glu Gly Leu Ser Ser Ile Lys Tyr Leu Phe	
	565 570	
	ggtgcttaat tgaaaaaaaa aaaaaaaaaa aaa	1762
25		
	<210> 9	
	<211> 25	
	<212> DNA	
	<213> Artificial Sequence	
30		
	<220>	
	<223> Designed oligonucleotide primer to obtain raffinose synthase gene.	
	<400> 9	
35	ccaatctgat catgcttgtg ccgaa	25
	<210> 10	
	<211> 30	
	<212> DNA	
40	<213> Artificial Sequence	

⟨223⟩ Designed oligonucleotide primer to obtain raffinose synthase gene.

5 <400> 10
ggaacaaagt tatgcactat tatttaaggt 30

10 <212> DNA
 <213> Artificial Sequence

15

<400> 11
ctaccaaat ccacaactta agttca 27

⟨213⟩ Artificial Sequence

25 <223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 12
ggaataataa gcttcacaca tactgtactc tc 32

30 <210> 13
 <211> 30
 <212> DNA
 <213> Artificial Sequence

35 <220>
<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

40 <400> 13
atggctccaa gcttttagcaa ggaaaattcc 30

<210> 14
<211> 30
<212> DNA
<213> Artificial Sequence
5
<220>
<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 14
10 tcaaaataag tactcaacag tggtaaaacc 30

<210> 15
<211> 30
<212> DNA
15 <213> Artificial Sequence

<220>
<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

20 <400> 15
ttggaagaga agacgccgcc gggaatcgtc 30

<210> 16
<211> 30
25 <212> DNA
<213> Artificial Sequence

<220>
<223> Designed oligonucleotide primer to obtain raffinose synthase gene.
30

<400> 16
ttaagccccg gcgagagctc tggccggaca 30

<210> 17
35 <211> 30
<212> DNA
<213> Artificial Sequence

<220>
40 <223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 17
accaatccaa aatctcatca aataatcgca 30

5 <210> 18
<211> 25
<212> DNA
<213> Artificial Sequence

10 <220>
<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 18
aaataatagg ggcagtacaa attacaccac 30

15 <210> 19
<211> 29
<212> DNA
<213> Artificial Sequence

20 <220>
<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 19
25 atggctccac cgagcgtaat taaatccga 29

<210> 20
<211> 30
<212> DNA
30 <213> Artificial Sequence

<220>
<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

35 <400> 20
ctaaaactca tacttaatag aagacaaacc 30

<210> 21
<211> 41
40 <212> DNA

⟨213⟩ Artificial Sequence

 $\langle 220 \rangle$

5 <223> Designed oligonucleotide primer to obtain raffinose synthase gene,
n is i.

<400> 21

cgatggatgg gnaanttnat ncancngan tggganatgt t 41

10 $\langle 210 \rangle$ 22

<211> 32

<212> DNA

⟨213⟩ Artificial Sequence

15 $\langle 220 \rangle$

<223> Designed oligonucleotide primer to obtain raffinose synthase gene,
n is i, r is a or g.

<400> 22

20 ggccacatnt tnacnarncc natnggngcn aa 32

<210> 23

<211> 30

<212> DNA

25 <213> Artificial Sequence

$\langle 220 \rangle$

<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

30 $\langle 400 \rangle$ 23

tgttactagg cgaaacaaga gtagctctga 30

<210> 24

<211> 47

35 $\langle 212 \rangle$ DNA

<213> Artificial Sequence

 $\langle 220 \rangle$

40 <223> Designed oligonucleotide primer to obtain raffinose synthase gene,
n is i.

	<400> 24	
	cgaggnggnt gncncncngg nttngtnatn atngangang gntggca	47
5	<210> 25	
	<211> 29	
	<212> DNA	
	<213> Artificial Sequence	
10	<220>	
	<223> Designed oligonucleotide primer to obtain raffinose synthase gene, n is i, y is t or c, r is a or g.	
	<400> 25	
15	atyttrtcna cngcnarrtc ytccatngt	29
	<210> 26	
	<211> 38	
	<212> DNA	
20	<213> Artificial Sequence	
	<220>	
	<223> Designed oligonucleotide primer to obtain raffinose synthase gene, n is i, y is t or c.	
25	<400> 26	
	ggnacntant ggytncangg ntgncanatg gtncantg	38
	<210> 27	
30	<211> 32	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
35	<223> Designed oligonucleotide primer to obtain raffinose synthase gene, n is i, r is a or g.	
	<400> 27	
	ggccacatnt tnacnarncc natngngncn aa	32
40		

<210> 28
<211> 26
<212> DNA
<213> Artificial Sequence
5
<220>
<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 28
10 atctatttgt catgacgatg atccga 26

<210> 29
<211> 30
<212> DNA
15 <213> Artificial Sequence

<220>
<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

20 <400> 29
ggccctcatt cccatattgg gatgacctc 30

<210> 30
<211> 30
25 <212> DNA
<213> Artificial Sequence

<220>
<223> Designed oligonucleotide primer to obtain raffinose synthase gene.
30
<400> 30

aagcatgcca aacatacaca tgctcaacag 30

35 <210> 31
<211> 30
<212> DNA
<213> Artificial Sequence

40 <220>

<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 31
agacccgggg aaagctttgg ggttactact 30

5

<210> 32
<211> 28
<212> DNA
<213> Artificial Sequence

10

<220>
<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 32
15 tggatgggaa actttataca ccctgact 28

<210> 33
<211> 28
<212> DNA
20 <213> Artificial Sequence

<220>
<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

25 <400> 33
gacatgttcc catctacaca cccttgatg 28

<210> 34
<211> 30
30 <212> DNA
<213> Artificial Sequence

<220>
<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

35 <400> 34
ccaatttatg ttagtgatgt tgttggcaag 30

<210> 35
40 <211> 26

	<212> DNA	
	<213> Artificial Sequence	
	<220>	
5	<223> Designed oligonucleotide primer to obtain raffinose synthase gene.	
	<400> 35	
	tcgactccca gggtagaatt gtcac	26
10	<210> 36	
	<211> 35	
	<212> DNA	
	<213> Artificial Sequence	
15	<220>	
	<223> Designed oligonucleotide primer to obtain raffinose synthase gene, n is i.	
	<400> 36	
20	cgattnaang tntggtggac nacncantgg gtngg	35
	<210> 37	
	<211> 38	
	<212> DNA	
25	<213> Artificial Sequence	
	<220>	
	<223> Designed oligonucleotide primer to obtain raffinose synthase gene, n is i, r is a or g.	
30	<400> 37	
	cantgnacca tntgncancc ntgnarccan tangtncc	38
	<210> 38	
35	<211> 30	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
40	<223> Designed oligonucleotide primer to obtain raffinose synthase gene.	

	<400> 38	
	gtaggggttc atatgaacac cttcaagctc	30
5	<210> 39	
	<211> 26	
	<212> DNA	
	<213> Artificial Sequence	
10	<220>	
	<223> Designed oligonucleotide primer to obtain raffinose synthase gene.	
	<400> 39	
	caacggcgag atcttgcatc gtcaac	26
15	<210> 40	
	<211> 30	
	<212> DNA	
	<213> Artificial Sequence	
20	<220>	
	<223> Designed oligonucleotide primer to obtain raffinose synthase gene.	
	<400> 40	
25	ggattcgaca caaaccgcca cgtcatcgtc	30
	<210> 41	
	<211> 27	
	<212> DNA	
30	<213> Artificial Sequence	
	<220>	
	<223> Designed oligonucleotide primer to obtain raffinose synthase gene.	
35	<400> 41	
	ccacgtgcac caccggaact tatcgac	27
	<210> 42	
	<211> 30	
40	<212> DNA	

⟨213⟩ Artificial Sequence

<220>

⟨223⟩ Designed oligonucleotide primer to obtain raffinose synthase gene.

5

<400> 42

aacatcgata ccatcggagt catgtccaat 30

<210> 43

10 $\langle 211 \rangle$ 30

<212> DNA

⟨213⟩ Artificial Sequence

<220>

15 <223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 43

gtagggttc atatgaacac cttcaagctc 30

20 $\langle 210 \rangle$ 44

<211> 29

<212> DNA

⟨213⟩ Artificial Sequence

25 $\langle 220 \rangle$

<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 44

tctacgtctg gcacgcgctt tgcggctac 29

30

<210> 45

<211> 31

<212> DNA

<213> Artificial Sequence

35

$\langle 220 \rangle$

<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 45

40 gttgacgtca tccacatatt ggagatgttg t 31

<223> Designed oligonucleotide linker to obtain raffinose synthase gene.

<400> 49
gatcgagctc gtgtcggatc cagct 25

5
<210> 50
<211> 17
<212> DNA
<213> Artificial Sequence

10
<220>
<223> Designed oligonucleotide linker to obtain raffinose synthase gene.

<400> 50
15 ggatccgaca cgagctc 17

<210> 51
<211> 30
<212> DNA
20 <213> Artificial Sequence

<220>
<223> Designed oligonucleotide primer to confirm direction of the inserted
raffinose synthase gene.

25
<400> 51
cctcctcgga ttccattgcc cagctatctg 30

<210> 52
30 <211> 30
<212> DNA
<213> Artificial Sequence

<220>
35 <223> Designed oligonucleotide primer to confirm direction of the inserted
raffinose synthase gene.

<400> 52
ggattcgaca caaaccgcca cgtcatcgtc 30

40

	<210> 53	
	<211> 29	
	<212> DNA	
	<213> Artificial Sequence	
5	<220>	
	<223> Designed oligonucleotide primer to confirm direction of the inserted raffinose synthase gene.	
10	<400> 53	
	tctacgtctg gcacgcgctt tgcggctac	29

What is claimed is:

1. A raffinose synthase gene which comprises a nucleotide sequence hybridizable with a nucleotide sequence selected from the group consisting of:

5 (a) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 1,

(b) the nucleotide sequence represented by SEQ ID NO: 2,

10 (c) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 3,

(d) the nucleotide sequence represented by the 236th to 2584th nucleotides in the nucleotide sequence represented by SEQ ID NO: 4,

15 (e) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 5,

(f) the nucleotide sequence represented by the 134th to 2467th nucleotides in the nucleotide sequence represented by SEQ ID NO: 6,

20 (g) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 7, and

(h) the nucleotide sequence represented by the 1st to 1719th nucleotides in the nucleotide sequence represented by SEQ ID NO: 8,
under stringent conditions, and encoding a protein being capable
25 of binding D-galactosyl group through α (1 \rightarrow 6) bond to the hydroxyl group attached to the carbon atom at 6-position of the D-glucose

residue in a sucrose molecule to form raffinose.

2. A raffinose synthase gene comprising a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 1.

5 3. A raffinose synthase gene comprising the nucleotide sequence represented by SEQ ID NO: 2.

4. A raffinose synthase gene comprising a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 3.

10 5. A raffinose synthase gene comprising the nucleotide sequence represented by the 236th to 2584th nucleotides in the nucleotide sequence represented by SEQ ID NO: 4.

15 6. A raffinose synthase gene comprising a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 5.

20 7. A raffinose synthase gene comprising the nucleotide sequence represented by the 134th to 2467th nucleotides in the nucleotide sequence represented by SEQ ID NO: 6.

8. A raffinose synthase gene comprising a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 7.

25 9. A raffinose synthase gene comprising the nucleotide sequence represented by the 1st to 1719th nucleotides in the nucleotide sequence represented by SEQ ID NO: 8.

10. A raffinose synthase gene comprising the nucleotide sequence represented by SEQ ID NO: 4, SEQ ID NO: 6 or SEQ ID NO: 8.

5 11. A nucleic acid comprising a partial nucleotide sequence of the raffinose synthase gene of any one of claims 1 to 10.

12. A method for detecting a nucleic acid containing a raffinose synthase gene which comprises detecting said nucleic acid by hybridization using the labeled nucleic acid of claim 10 11 as a probe.

13. A method for amplifying a nucleic acid containing a raffinose synthase gene which comprises amplifying said nucleic acid by polymerase chain reaction (PCR) using the nucleic acid of claim 11 as a primer.

15 14. A method for obtaining a raffinose synthase gene which comprises the steps of:

detecting a nucleic acid containing said raffinose synthase gene by hybridization using the labeled nucleic acid of claim 11 as a probe, and

20 recovering the detected nucleic acid.

15. A method for obtaining a raffinose synthase gene which comprises the steps of:

amplifying a nucleic acid containing said raffinose synthase gene by PCR using the nucleic acid of claim 11 as a 25 primer, and

recovering the amplified nucleic acid.

16. A nucleic acid comprising a nucleic acid containing the raffinose synthase gene of any one of claims 1 to 10 which is joined to a nucleic acid exhibiting promoter activity in a host cell.

5 17. A vector comprising the raffinose synthase gene of any one of claims 1 to 10.

18. A transformant, wherein the raffinose synthase gene of any one of claims 1 to 10 is introduced into a host cell.

10 19. A transformant, wherein the nucleic acid of claim 16 is introduced into a host cell.

20. A transformant, wherein the vector of claim 17 is introduced into a host cell.

15 21. The transformant of any one of claims 18 to 20, wherein the host is a microorganism.

22. The transformant of any one of claims 18 to 20, wherein the host is a plant.

23. A method for producing a raffinose synthase which comprises the steps of:

20 culturing or growing the transformant of any one of claims 18 to 22 to produce the raffinose synthase, and collecting the raffinose synthase.

24. A raffinose synthase comprising the amino acid sequence represented by SEQ ID NO: 1.

25 25. A raffinose synthase comprising the amino acid sequence represented by SEQ ID NO: 3.

26. A raffinose synthase comprising the amino acid sequence represented by SEQ ID NO: 5.

27. A raffinose synthase comprising the amino acid sequence represented by SEQ ID NO: 7.

Abstract of the disclosure:

A raffinose synthase gene comprising a nucleotide sequence hybridizable with a nucleotide sequence selected from the group consisting of: (a) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 1, (b) the nucleotide sequence represented by SEQ ID NO: 2, (c) a nucleotide sequence encoding the amino acid sequence of represented by SEQ ID NO: 3, (d) the nucleotide sequence represented by SEQ ID NO: 4 or by the 236th to 2584th nucleotides in the nucleotide sequence represented by SEQ ID NO: 4, (e) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 5, (f) the nucleotide sequence represented by SEQ ID NO: 6 or by the 134th to 2467th nucleotides in the nucleotide sequence represented by SEQ ID NO: 6, (g) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 7, and (h) the nucleotide sequence represented by SEQ ID NO: 8 or by the 1st to 1719th nucleotides in the nucleotide sequence represented by SEQ ID NO: 8, under stringent conditions, and encoding a protein being capable of binding D-galactosyl group through α (1 \rightarrow 6) bond to the hydroxyl group attached to the carbon atom at 6-position of the D-glucose residue in a sucrose molecule to form raffinose is disclosed.